

Mark Scheme (Results)

Summer 2021

Pearson Edexcel International Advance Level In Biology (WBI12) Paper 01 Cells, Development, Biodiversity and Conservation

| Question | Answer | Mark |
|----------|---|------|
| Number | | |
| 1(a)(i) | | |
| | The only correct answer is B β-glucose | |
| | A is not correct because the molecule is β-glucose | |
| | | |
| | C is not correct because the molecule is β-glucose | |
| | D is not correct because the molecule is β -glucose | (1) |

| Question | Answer | Mark |
|----------|---|------|
| Number | | |
| 1(a)(ii) | The only correct answer is C form microfibrils | |
| | A is not correct because cellulose forms microfibrils | |
| | B is not correct because cellulose forms microfibrils | |
| | D is not correct because cellulose forms microfibrils | (1) |

| Question | Answer | Mark |
|-----------|--|------|
| Number | | |
| 1(a)(iii) | The only correct answer is B chlorophyll A is not correct because magnesium is not a component of calcium pectate C is not correct because magnesium is not a component of DNA | |
| | D is not correct because magnesium is not a component of starch | (1) |

| Question | Answer | Additional guidance | Mark |
|----------|---------|---------------------|------|
| Number | | | |
| 1(b)(i) | | | |
| | • xylem | | (1) |
| | | | |

| Question Number | Answer | Additional guidance | Mark |
|--------------------|----------------|---------------------|------|
| 1(b)(ii) | • sclerenchyma | Accept tracheids | (1) |

| Question | Answer | Additional guidance | Mark |
|----------|--|---|------|
| Number | | | |
| 2(a)(i) | A calculation in which: | Example of calculation: | |
| | conversion of actual length of scale into nm (1) actual length divided by the given length (answer given to 2 significant figures and in standard form) (1) | 35mm = 35000 000 nm 35000 000 ÷1200 = ×2.9 x 10 ⁴ accept +/- 1mm for length of bar | (2) |

| Question | Answer | Additional guidance | Mark |
|----------|---|--|------|
| Number | | | |
| 2(a)(ii) | An explanation that makes reference to two of the following points: | ecf from 2(a)(i) applies | |
| | | | |
| | electron microscope (1) | | |
| | due to high magnification (1) | Accept high resolution | |
| | duc to high magnification (1) | Accept converse for light microscope | (2) |
| | | Accept 3D image if qualified as scanning | (2) |
| | | electron microscope | |

| Question | Answer | Additional guidance | Mark |
|----------|--|---------------------|------|
| Number | | | |
| 2(b)(i) | An answer that makes reference to the following points: | | |
| | presence of {(peptidoglycan) cell wall / circular chromosome(s) / DNA associated with histones / 70S ribosomes / RNA polymerase} (1) | Accept ether bonds | (2) |
| | absence of {nucleus / membrane bound organelles} (1) | | (2) |

| Question | Answer | Additional guidance | Mark |
|----------|---|---|------|
| Number | | | |
| 2(b)(ii) | An explanation that makes reference to three of the following points: | | |
| | (analysis of) molecular evidence (1) | e.g. DNA, mRNA, proteins, enzymes | |
| | to identify similarities and differences in biological molecules / comparison of biological molecules (1) | | |
| | (analysis of) phenotype (1) | e.g. cell structure / anatomical features, grow in different {habitats/ conditions} | |
| | to identify similarities and differences between the two microorganisms (phenotype) (1) | e.g. named similarities and differences in cell structure | (3) |

| Question | Answer | Mark |
|----------|---|------|
| Number | | |
| 3(a)(i) | | |
| | The only correct answer is C two | |
| | | |
| | A is not correct because there is one maternal and one paternal chromosome with three gene loci | |
| | | |
| | B is not correct because there is one maternal and one paternal chromosome with three gene loci | |
| | | (4) |
| | D is not correct because there is one maternal and one paternal chromosome with three gene loci | (1) |
| | | |

| Question | Answer | Additional guidance | Mark |
|----------|--|--|------|
| Number | | | |
| 3(a)(ii) | A description that makes reference to the following points: | | |
| | uncoiling (of chromosome) (1) | Accept correct reference to histones / to form chromatin | |
| | • used in protein synthesis / DNA replication (in S phase) (1) | Accept transcription | (2) |

| Question | Answer | Additional guidance | Mark |
|----------|------------|---------------------|------|
| Number | | | |
| 3(b)(i) | | | |
| | prophase I | | (1) |
| | | | |

| Question | Answer | Mark |
|----------|---|------|
| Number | Additional guidance | |
| 3(b)(ii) | | |
| | The only correct answer is C three | |
| | | |
| | A is not correct because ABe could not be formed by one chiasmata | |
| | | |
| | B is not correct because ABe could not be formed by one chiasmata | |
| | | |
| | D is not correct because ABe could not be formed by one chiasmata | (1) |
| | | |

| Question | Answer | Additional guidance | Mark |
|-----------|---|-----------------------------|------|
| Number | | | |
| 3(b)(iii) | An explanation that makes reference to two of the following points: | | |
| | | | |
| | alleles b and e are closer together (1) | Accept converse for A and E | |
| | | | |
| | therefore {crossing over / chiasma} is less likely to occur | Accept converse for A and E | (2) |
| | between these alleles (1) | | |
| | | | |

| Question | Answer | Additional guidance | Mark |
|----------|--|---------------------|------|
| Number | | | |
| 4(a)(i) | A description that makes reference to the following points: | | |
| | the germination of the seeds remains {high/ constant} for an initial period of time (1) | | |
| | (after this) the germination of the seeds (rapidly) {decreases / reduces} with increased time in storage (1) | | (2) |

| Question | Answer | Additional guidance | Mark |
|----------|--|--|------|
| Number | | | |
| 4(a)(ii) | An explanation that makes reference to four of the following points: | | |
| | (many different plants result in) increased genetic diversity of the seeds stored (1) | | |
| | because more (seeds from different) {individuals/ plants} increases the probability of having {more alleles / different genotypes / heterozygotes} (1) | Accept reduces chance of seeds from (mainly) {homozygous recessive / diseased} plants | |
| | adult plants allow {pollination / sexual reproduction} to occur (1) | | |
| | to produce new seeds (for storage) (1) | Accept to produce seeds which are genetically different (from parent plants) | |
| | to replace those seeds stored for a long time (1) | Accept to increase the {viability of stored seeds / percentage of seeds that will germinate} | (4) |

| Question | Answer | Additional guidance | Mark |
|----------|---|--|------|
| Number | | | |
| 4(b)(i) | A calculation in which: | Example of calculation | |
| | mass of cutlery per month calculated (1) mass of cutlery per year calculated to two significant figures with | $(130000 \div 100) \times 60 = 78000 \text{ (kg)}$ $(78000 \times 12) = 940000 \text{ kg} / 9.4 \times 10^5 \text{ kg}$ | |
| | unit (1) | Correct answer with no working scores full marks | (2) |

| Question | Answer | Additional guidance | Mark |
|----------|--|---|------|
| Number | | | |
| 4(b)(ii) | An answer that includes the following points:more (cutlery) can (constantly) be made from avocado seeds | Accept carbon neutral / lower | |
| | (1) | carbon emissions (than oil-based plastic production) / biodegradable Accept as avocado seeds are waste products | |
| | as avocados are {continually being grown / can be grown by future generations} (1) | | (2) |

| Question Number | Answer | Additional guidance | Mark |
|--------------------|---|----------------------------------|------|
| 5(a)(i) | An answer that includes the following points: | Example of diagram | |
| | at least two curved cisternae drawn (without ribosomes or interconnections) (1) | cisternae Cisternae | |
| | vesicle drawn (1) | Cisternae | |
| | cisternae labelled (1) | | |
| | second correct label (1) | trans face — (secretory) vesicle | |
| | | | (4) |

| Question | Answer | Mark |
|----------|---|------|
| Number | | |
| 5(a)(ii) | The only correct answer is C two | |
| | A is not correct because the Golgi apparatus is involved in the formation of extracellular enzymes and modification of proteins | |
| | B is not correct because the Golgi apparatus is involved in the formation of extracellular enzymes and modification of proteins | |
| | D is not correct because the Golgi apparatus does not form peptide bonds | (1) |

| Question | Answer | Mark |
|-----------|---|------|
| Number | | |
| 5(a)(iii) | | |
| | The only correct answer is D ribosomes | |
| | | |
| | A is not correct because the ribosomes would become radioactive first | |
| | | |
| | B is not correct because the ribosomes would become radioactive first | |
| | D is not so west because the ribesomes would be some radio active first | (4) |
| | D is not correct because the ribosomes would become radioactive first | (1) |
| | | |

| Question | Answer | Additional guidance | Mark |
|----------|------------------------------------|-----------------------------|------|
| Number | | | |
| 5(b)(i) | | Example of calculation | |
| | • 120(%) / 54.5(%) / 54(%) / 55(%) | $\frac{0.5-1.1}{0.5}$ x 100 | (1) |

| Question | Answer | Mark |
|----------|---|------|
| Number | | |
| 5(b)(ii) | | |
| | The only correct answer is B interphase | |
| | A is not correct because the Golgi apparatus would increase in size in interphase | |
| | C is not correct because the Golgi apparatus would increase in size in interphase | |
| | D is not correct because the Golgi apparatus would increase in size in interphase | (1) |

| Question | Answer | Additional guidance | Mark |
|-----------|---|---|------|
| Number | | | |
| 5(b)(iii) | An answer that makes reference to four of the following points: | | |
| | mitosis / two cells formed from one cell (1) | | |
| | therefore {number / size} of Golgi (in each cell) has to increase (in interphase / G1) (1) | | |
| | to provide enough cell {contents / organelles} for two (daughter) cells (after mitosis has occurred) (1) | | |
| | because increased {protein synthesis / protein modification / gene expression} will occur (in interphase) (1) | Accept relationship between {number/size} of Golgi and level of protein synthesis | |
| | because named protein(s) are required (by the cell) (1) | e.g. enzymes, hormones, structural proteins | (4) |

| Question Number | Answer | Additional guidance | Mark |
|--------------------|---|----------------------------|------|
| 6(a) | An explanation that includes four of the following points: | | |
| | cortical granules are released (from the egg cell) (1) | | |
| | cortical granules fuse with the {zona pellucida / egg cell membrane} (1) | | |
| | cortical granules release {chemicals / enzymes / proteins} to cause the hardening of zona pellucida (1) | | |
| | prevents more than one sperm (cell) entering the egg cell (1) | ACCEPT prevents polyspermy | (4) |
| | egg cell and sperm nuclei are haploid and they fuse (during fertilisation) (1) | | |

| Question Number | Answer |
|--------------------|---|
| = | Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlined in the generic mark scheme. The indicative content below is not prescriptive and candidates are not required to include all the material indicated as relevant. Additional content included in the response must be scientific and relevant. Iength of middle region |
| | (6 |

| | | | Additional guidance |
|---------|-----|--|---|
| Level 0 | 0 | No awardable content | |
| Level 1 | 1-2 | Limited number of the most important or relevant scientific factors from the data/information provided are synthesised. No judgement is made. | 1 mark – either R or G for one mating strategy 2 marks – either R+G for one mating strategy OR either R or G for two mating strategies OR 1 mark - basic information (I) given 2 marks - basic information (I) given + 1 or 2 x {R or G} |
| Level 2 | 3-4 | Some of the most important or relevant scientific factors from the data/information provided are synthesised. A limited accurate judgement is made. | 3 marks – 3 x (R or G) considered 4 marks = 4 x (R or G) considered |
| Level 3 | 5-6 | Most of the important or relevant scientific factors from the data/information provided are synthesised. A detailed and accurate judgement is made. | 5 marks = either reproductive success for all strategies considered plus genetic diversity of two strategies considered OR genetic diversity for all strategies considered plus reproductive success of two strategies considered 6 marks genetic diversity and reproductive success considered for all strategies |

| Question | Answer | Additional guidance | Mark |
|----------|--|--|------|
| Number | | | |
| 7(a) | A calculation in which: | Example of calculation | |
| | total population of Santa Cruz in 2020 calculated (1) total population of Santa Cruz in 2025 calculated (1) | $(17000 \div 85) \times 100 = 20000$ $20000 \times 1.064^5 = 27273.3 / 27273.328 / 27273.33$ | |
| | population given to whole number (1) | {27 273 / 27 000} Correct answer with no working scores full marks ecf applies {26400 / 26000} = 2 marks | (3) |

| Question | Answer | Additional guidance | Mark |
|----------|---|---|------|
| Number | | | |
| 7(b) | positive correlation between the number of recorded non endemic species and the number of tourists increased | ACCEPT number of recorded non endemic species increased as the number of tourists increased | (1) |

| Question Number | Answer | Additiona | l guidance | | | Mark |
|--------------------|--|------------------------|---------------------------|-------------|----------------|------|
| 7(c)(i) | A calculation in which: | Example of calculation | | | | |
| | N(N-1) correctly calculated (1) | (147 x 146 | 6) = 21462 | | | |
| | ∑n(n-1) correctly calculated (1) | 4684 | | | | |
| | • correct substitution to obtain D to 1 d.p. (1) | 4.6 (4.58 = 2 i | marks) | | | |
| | | Species | Number of individuals (n) | (n-1) | n(n-1) | |
| | | А | 21 | 20 | 420 | |
| | | В | 2 | 1 | 2 | |
| | | С | 4 | 3 | 12 | |
| | | D | 13 | 12 | 156 | |
| | | Е | 54 | 53 | 2862 | |
| | | F | 15 | 14 | 210 | |
| | | G | 6 | 5 | 30 | |
| | | Н | 32 | 31 | 992 | |
| | | | Total (N)=147 | | ∑n(n-1)=4684 | |
| | | Correct a | | working sco | res full marks | (3) |

| Question | Answer | Additional guidance | Mark |
|-----------------|--|--|------|
| Number 7(c)(ii) | An answer that includes five of the following points: | | |
| | • reduction in biodiversity (1) | Accept possible increase in species that have blackberry as part of food chain | |
| | as forest is habitat for many species of plants (1) | Accept reduction in forest habitat (for many species unqualified) / blackberry {outcompetes (the forest) plants / introduces disease} | |
| | (because) populations will decrease because loss of forest will result in {reduced habitat for animals / reduced food} / increased competition between (animal) species} (1) | Accept animals not adapted to feed on blackberry / blackberry is poisonous to the animals Accept descriptions of what animals would compete for e.g. nesting sites | |
| | causing reduction in species richness (1) | | |
| | (because) loss of forest will result in reduction of genetic diversity (1) | Accept reduction in gene pool | (5) |
| | • (due to) reproduction in {smaller / isolated} populations (1) | | (5) |

| Question Number | Answer | Additional guidance | Mark |
|--------------------|--|--|------|
| 8(a) | An answer that includes at least one similarity and one difference: | | |
| | similarities: | | |
| | both contain membrane bound organelles (1) | ACCEPT named organelle(s) e.g. nucleus, mitochondria, 80S ribosomes etc ACCEPT both contain DNA associated with histones (1) | |
| | both contain a {cell membrane / cell wall} (1) | | |
| | max 2 differences: | | |
| | plant cell wall is composed of cellulose whereas a fungal cell wall is composed of chitin (1) | | |
| | fungal cells contain glycogen granules whereas plant cells contain {starch grains / amyloplasts} (1) | | |
| | plant cells contain {chloroplasts / plasmodesmata / tonoplast} whereas fungal cells do not (1) | ACCEPT fungal cells contain lysosomes whereas plant cells do not / plant vacuole may be larger than fungal vacuole | (3) |

| Question | Answer | Additional guidance | Mark |
|----------|---|---|------|
| Number | | | |
| 8(b)(i) | An explanation that makes reference to four of the following points: | | |
| | • reduction in growth (1) | | |
| | and any three from: | | |
| | | | |
| | due to reduced {cell division / mitosis} (1) | ACCEPT reduced DNA synthesis results in fewer cells in S phase / not enough DNA produced for mitosis to occur | |
| | reduced gene expression will cause reduction in {enzyme/ protein} production (1) | ACCEPT named enzyme e.g. DNA polymerase | |
| | reduced phospholipids which are needed for (growth of) {cell / organelle} membranes (in daughter cells) (1) | | |
| | reduced transport of sucrose (around the plant) in the phloem / reduced sucrose available for other plant cells (1) | ACCEPT correct examples of sucrose use by plant e.g. reduced {respiration / ATP production} | (4) |

| Question Number | Answer |
|--------------------|--|
| *8(b)(ii) | Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlined in the generic mark scheme. The indicative content below is not prescriptive and candidates are not required to include all the material indicated as relevant. Additional content included in the response must be scientific and relevant. |
| | Description of data infection with fungus results in increased {tannin/ flavonoid} production increasing the concentration of tannin reduces the leaf damage from herbivores negative correlation / linear decrease increasing the concentration of flavonoids increases the diameter of inhibition zone doubling the concentration of flavonoids increases the diameter of inhibition zone by 28.6% Plant |
| | explanation of how tannins reduce leaf damage e.g. herbivores don't like the taste / toxic / poisonous therefore less likely to eat the plant (A) more leaves for photosynthesis and therefore increased growth (A) explanation of what the diameter of inhibition zone shows = link to bactericidal/ bacteriostatic effect of flavonoids (A) reduced plant infections (A) link to why reduced herbivory and reduced plant infection benefits the infected plant e.g. fewer resources goes into growing new leaves and more can go into sexual reproduction (A) disadvantage for plant – increased fungal growth (D) more resources required {to produce tannins / for gene expression} instead of being used for {growth / sexual |
| | more resources required (to produce tannins / for gene expression) instead of being used for (growth / sexual reproduction) (D) Fungus link to why reduced herbivory benefits the fungus e.g. reduced competition for plant resources from herbivores, more plant resources available to fungus (A) link to why reduced plant infection benefits the fungus e.g. reduced competition for plant resources from {bacteria / other fungi} (A) reduced resources for fungus from plant as more used to synthesise tannins / flavonoids (D) |

| | the fungus might only be able to {influence / live in / complete life cycle in} darnel plants (D) | | |
|---------|---|--|---|
| | | | Additional guidance |
| Level 0 | 0 | No awardable content | |
| Level 1 | 1-2 | Demonstrates isolated elements of biological knowledge and understanding to the given context with generalised comments made. Vague statements related to consequences are made with limited linkage to a range of scientific ideas, processes, techniques and procedures. The discussion will contain basic information with some attempt made to link knowledge and understanding to the given context. | Information from one section |
| Level 2 | 3-4 | Demonstrates adequate knowledge and understanding by selecting and applying some relevant biological facts/concepts. Consequences are discussed which are occasionally supported through linkage to a range of scientific ideas, processes, techniques and procedures. The discussion shows some linkages and lines of scientific reasoning with some structure. | level one content plus consideration of advantage(s) and / or disadvantage(s) 3 marks = L1+ one from {FA / FD / PA / PD} 4 marks = L1+ two from {FA / FD / PA / PD} FA = fungal advantage FD = fungal disadvantage PA = plant advantage PD = plant disadvantage |
| Level 3 | 5-6 | Demonstrates comprehensive knowledge and understanding by selecting and applying relevant knowledge of biological facts/concepts. Consequences are discussed which are supported throughout by sustained linkage to a range of scientific ideas, processes, techniques and procedures. The discussion shows a well-developed and sustained line of scientific reasoning which is clear and logically structured. | Information from three sections applied to the given context showing a good understanding of the advantages and disadvantages to the infected plant and to the fungus 5 marks = L2 + three from {FA / FD / PA / PD} 6 marks - all four of {FA / FD / PA / PD} discussed |

| Question | Answer | Additional guidance | Mark |
|-----------|---|--|------|
| Number | | | |
| 8(b)(iii) | A description that includes reference to the following points: | | |
| | (flavonoid drug) tested on small numbers of patients with {P. aeruginosa / lung infection} (1) use of double-blind trial (1) | ACCEPT tested on {100 to 500} of patients with { <i>P. aeruginosa /</i> lung infection} ACCEPT placebo / established drug treatment | |
| | | | (2) |
| | | | |