



Mark Scheme (Results)

June 2022

Pearson Edexcel International Advanced Level
In Biology (WBI13) Paper 01
Practical Skills in Biology I

Question Number	Answer	Additional Guidance	Mark
1(a)(i)	An explanation that includes 3 of the following points: • it {carries / transports} the male {gametes / nuclei / generative nucleus} (1) • to the ovule / micropyle / ovary / ovum / female gamete (1) • to {fuse with / fertilise} {ovum / egg (cell) / female gamete / female nucleus / polar nuclei} (1) • digest tissue of style (1)	Accept sperm	(3)

Question Number	Answer	Additional Guidance	Mark
1(a)(ii)	An explanation that includes the following points: • the tubes grow towards the {micropyle / ovule / ovary / ovum / female gamete} (1) • {due to chemicals (released by the embryo sac) / it is chemotropic} (1)		(2)

Question Number	Answer	Additional Guidance	Mark
1(b)(i)	An answer that includes the following: • percentage germination / number germinating (1) • pollen tube length (1)		(2)

Question Number	Answer	Additional Guidance	Mark
1(b)(ii)	An answer that includes the following points: • it involves {enzymes / (chemical) reactions} (1) • example of how {enzymes / (chemical) reactions} are affected by temperature (1)	e.g kinetic energy / denaturation	(2)

Question Number	Answer	Additional Guidance	Mark
1(c) (i)	An answer that includes the following points: • description of method (1) • calculation of dilution factor / use of $C_1V_1=C_2V_2$ to calculate sucrose solution volume as 2 (cm³) (1) • stating the volume of sucrose solution and calcium ion solution to be used to make 10cm³ (1)	ACCEPT mixed / dilute 5x / by 5/ 1:4 / 1 in 5 e. g. 2 cm³ of sucrose solution to 8 cm³ calcium ion solution / allow removal of 10cm³ of bigger volume allow ecf	(3)


Question Number	Answer	Additional Guidance	Mark
1(c)(ii)	varying the (concentration of) calcium ion solution (1)	ACCEPT use of buffer (solution)	(1)

Question Number	Answer	Additional Guidance		Mark												
1(d)(i)	An answer including the following: • suitable table drawn (1) • all headings correct with units (1) • all data for germination (only) and the 5 sucrose concentrations entered correctly (1)	<table><tr><th>Sucrose concentration / solution / mol dm⁻³</th><th>Germination (%)</th></tr><tr><td>0</td><td>6</td></tr><tr><td>0.2</td><td>46</td></tr><tr><td>0.4</td><td>70</td></tr><tr><td>0.8</td><td>23</td></tr><tr><td>1.6</td><td>0</td></tr></table>	Sucrose concentration / solution / mol dm ⁻³	Germination (%)	0	6	0.2	46	0.4	70	0.8	23	1.6	0	Allow ± 0.5 if they choose to quote germination % to 1 dp.	(3)
Sucrose concentration / solution / mol dm ⁻³	Germination (%)															
0	6															
0.2	46															
0.4	70															
0.8	23															
1.6	0															

Question Number	Answer	Additional Guidance	Mark
1(d)(ii)	An answer that includes the following points : • both (appear to) have optimum <u>at 0.4 mol dm⁻³</u> (1) • (but for either) it could be anywhere between above 0.2 mol dm⁻³ and below 0.8 mol dm⁻³ (1) • values for optimum could be different (to each other) / they might not have the same optima / one optimum might not be 0.4 mol dm⁻³ (1) • more concentrations of sucrose (between 0.2 and 0.8 mol dm⁻³) should be investigated (1)	ACCEPT 0.2 - 0.8 mol dm ⁻³	(3)

Question Number	Answer	Additional Guidance	Mark
2(a)	An answer that includes the following points : • cut root tip • (root tips) placed in (warm) acid (1) • (root tips) then placed in named stain (1) • (root tip placed on a microscope slide and) {macerated / teased / described} / squashed (1) • use of high power (on a microscope) (1) • correct ref to safety issue (1)	e.g. (acetic / ethanoic) orcein / toluidine blue / methylene blue e.g. rinsing tips in water / wearing gloves / goggles	(5)

Question Number	Answer	Additional Guidance	Mark
2(b)	An answer that includes the following steps: - correct total cell count and total dividing cell count (1) - division of dividing cell count by total cell count (and multiply by 100) (1)	NO ecf e.g. $40 / 4$ ACCEPT 3 ACCEPT any figure in range 35 to 46 e.g. $(4 \div 40) = 0.10$ ($\times 100 = 10.0 (\%)$)	(2)

Question Number	Answer	Additional Guidance	Mark												
2(c)(i)	A graph with the following features : - A axes correct (x - appropriate to their graph , y - mitotic index) and y axis with no break in the axis (1) - L axes correctly labelled with units (1) - P correct plotting on a linear scale on y (1) - S bar chart (1)	e.g.  <table><caption>Mitotic Index Data</caption><thead><tr><th>Treatment</th><th>Mitotic Index</th></tr></thead><tbody><tr><td>A 1 (mg cm⁻³)</td><td>4.8</td></tr><tr><td>A 7 (mg cm⁻³)</td><td>1.2</td></tr><tr><td>B 1 (mg cm⁻³)</td><td>4.2</td></tr><tr><td>B 7 (mg cm⁻³)</td><td>1.8</td></tr><tr><td>Control (mg cm⁻³)</td><td>7.5</td></tr></tbody></table>	Treatment	Mitotic Index	A 1 (mg cm ⁻³)	4.8	A 7 (mg cm ⁻³)	1.2	B 1 (mg cm ⁻³)	4.2	B 7 (mg cm ⁻³)	1.8	Control (mg cm ⁻³)	7.5	(4)
Treatment	Mitotic Index														
A 1 (mg cm ⁻³)	4.8														
A 7 (mg cm ⁻³)	1.2														
B 1 (mg cm ⁻³)	4.2														
B 7 (mg cm ⁻³)	1.8														
Control (mg cm ⁻³)	7.5														

Question Number	Answer	Additional Guidance	Mark
2(c)(ii)	An answer that includes the following points : • each treatment should be repeated (1) • {all conditions / named condition} should be kept constant (1) • (mean and) SD calculated / error bars (1) • look for overlap in SDs / perform t-test (1)	IGNORE standard conditions ALLOW range bars as ecf	(4)

Question Number	Answer	Additional Guidance	Mark
3(a)	An answer that includes the following points : • serial dilution of stock solution / described (1) • describe method to measure out same volume of each dilution (into test tube) (1) • add same {volume / no. of drops} (of same concentration) iodine (solution to each test tube) (1) • description of method for observing colour of solution (1) • {observe / note down} the colour of the resultant solution (1)	e.g. putting solution in a test tube of the same diameter	(4)

Question Number	Answer	Additional Guidance	Mark
3(b)	An answer that includes the following points: • Russet and King Edward show {high(er) level of starch / black and dark blue} and therefore best for baking (1) • Nicola and Purple Congo show {low(er) level of starch / pale blue } and therefore best for boiling (1) • King Edward better than Russet for baking / Nicola better than Purple Congo for boiling / King Edward best for baking, Nicola best for boiling(1) • therefore the suggestions are supported by the data (1) • a comment on the subjectivity of the colour differences (1)	e.g. the results are semi-quantitative	(4)

Question Number	Answer	Additional Guidance	Mark
3(c)	An answer including the following points : • suitable method chosen (1) • further detail (1)	e.g colorimeter, looking down on test tubes and adjusting heights e.g. colorimeter, zeroing, use of same cuvette, measure absorbance or transmission looking down tubes, judging colour intensity by looking down and measure heights.	(2)

Question Number	Answer	Additional Guidance	Mark
3(d)(i)	An answer including the following steps : • correct reading at day 23 minus correct reading at day 15 and correct subtraction(1) • division of answer by time (8 days) (1)	e.g. $14-5 = 9$ or $5-14 = -9$ e.g. $9 \div 8 = 1.125 \text{ au day}^{-1}$ ACCEPT au / day, au per day ACCEPT 1.1, 1.13 IGNORE the sign	(2)

Question Number	Answer	Additional Guidance	Mark
3(d)(ii)	An answer including the following points : • negative correlation / falls throughout / non-linear (1) • falls {very little / slowly / by 2 au} {between / up to} days 0 and {5 / 15} • falls {a lot / rapidly / by 9} {between / up to} days 15 and 23 (1)		(2)

Question Number	Answer	Additional Guidance	Mark
3(d)(iii)	A graph that shows the following features: • overall rise (1) • an inflection at 15 and then steeper than anywhere else on their graph (1)	soluble sugar content / au days in storage	(2)