




Mark Scheme (Results)

Summer 2024

Pearson Edexcel International Advanced
Subsidiary Level In Chemistry (WCH13)
Paper 01R: Practical Skills in Chemistry I

Question Number	Answer	Additional Guidance	Mark
1(a)	• diagram of a gas syringe	<u>Example of a diagram:</u>  0.150 mol dm⁻³ sulfuric acid and 0.200 g sodium sulfite Gas syringe should have plunger and graduations and be connected to the delivery tube.	(1)

Question Number	Answer	Additional Guidance	Mark
1(b)	• calculation of M_r of Na₂SO₃ (1) • calculation of number of moles (1)	Example of a calculation: $((23.0 \times 2) + 32.1 + (3 \times 16) =) 126.1$ Allow 126 $0.200 \div 126.1 = 1.5860 \times 10^{-3} / 0.0015860$ Allow $1.5873 \times 10^{-3} / 0.0015873$ TE from M1 to M2 Ignore SF except 1SF on M2 Correct answer scores (2)	(2)

Question Number	Answer	Additional Guidance	Mark
1(c)	- calculation of gas volume (1) - conversion to dm³ (1)	<u>Example of a calculation:</u> $29.5 / 0.001586 = 18\,600 \text{ (cm}^3\text{)}$ $18\,600 \div 1000 = 18.6 \text{ dm}^3 / 19 \text{ dm}^3$ Alternate method: Convert to dm³ $0.0295 / 2.95 \times 10^{-2}$ Calcⁿ of gas vol $0.0295 \div \text{mol} = 18.6$ TE from (b) Allow litres in place of dm³ Ignore SF except 1SF Ignore incorrect standard form Ignore dm³ mol⁻¹	(2)

Question Number	Answer	Additional Guidance	Mark
1(d)(i)	use of $n \div c = v$ with units	<u>Example of a calculation:</u> $0.00159 \div 0.150 = 0.010573 / 0.0106 \text{ dm}^3 / 10.6 \text{ cm}^3$ TE from (b) Allow litres in place of dm³ Ignore SF except 1SF	(1)

Question Number	Answer	Additional Guidance	Mark
1(d)(ii)	suitable suggestion	e.g. temperature is lower than standard conditions, some gas escapes before the bung can be replaced, some gas escapes as the reaction starts, SO₂ dissolved in the sulfuric acid Allow dissolved in sulfuric acid / sodium sulfate solution Ignore incomplete reaction Ignore impure reactants Ignore measurement error Ignore solubility in water / reacts with water Ignore “not standard conditions” Ignore references to pressure	(1)

Question Number	Answer	Additional Guidance	Mark
1(e)	An answer that makes reference to the following point: sulfur dioxide / SO₂ is toxic	Allow poisonous Allow irritant Allow choking smell Ignore acid rain / pollution / breathing difficulties Do not award “toxic” without named gas	(1)

(Total for Question 1 = 8 marks)

Question Number	Answer	Additional Guidance	Mark
2(a)(i)	A description that makes reference to the following points: - Li fizzes / moves (more slowly on the surface of the water) - sodium fizzes more / moves faster or forms a sphere on the water - potassium produces a lilac flame	(1) Allow floats Ignore smoke Do not award “produces precipitate” Sinks to bottom or flame negates M1 Do not award M1 if Lithium is stated as the most reactive (1) Allow yellow(-orange) flame produced for sphere (1) Allow purple for lilac Allow always produces a flame (no colour required) Ignore explosion If no other mark scored allow “Reactivity increases from Li to K” (or description) or “metal disappears/dissolves more quickly from Li to K” (or converse) for 1 mark	(3)

Question Number	Answer	Additional Guidance	Mark
2(a)(ii)	An answer that makes reference to two of the following points: • use small pieces of metal (1) • use tongs / gloves (to move the metal) (1) • ensure all oil is removed from the surface of the metal (1) • use a large container / volume of water (1)	Allow use less solid Do not award “wear gloves” if related to heat Allow use a safety screen or fume hood Allow lab coat Ignore distance / open space Ignore presence of flames Ignore masks	(2)

Question Number	Answer	Additional Guidance	Mark
2(b)(i)	An answer that makes reference to the following points: • balanced ionic equation and state symbols	$\text{Mg}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Mg}(\text{OH})_2(\text{s})$	(1)

Question Number	Answer	Additional Guidance	Mark
2(b)(ii)	An answer that makes reference to the following point: • white and precipitate	Allow ppt/ppte/solid/suspension Allow goes cloudy/milky	(1)

Question Number	Answer	Additional Guidance	Mark
2(b)(iii)	A description that makes reference to the following points: • add (dilute) sulfuric acid / aqueous sulfate ions (1) • a white precipitate (of barium sulfate / BaSO₄) (1)	Accept named solutions Ignore additional acids Do not award concentrated sulfuric acid Allow ppt/ppte/solid M2 dependent on M1 or a near-miss (e.g. incorrect formula) Ignore further tests	(2)

Question Number	Answer	Additional Guidance	Mark
2(b)(iv)	An answer that makes reference to the following points: • magnesium: no colour (1) • barium: (apple) green (1)	Allow colourless Do not award white Do not award blue-green	(2)

(Total for Question 2 = 11 marks)

Question Number	Answer	Additional Guidance	Mark
3(a)	An answer that makes reference to the following points: • colourless • (to) pink	(1) (1) Do not award purple / red 1 mark if colours are correct but inverted	(2)

Question Number	Answer	Additional Guidance	Mark
3(b)	• mean titre calculated	Example of a calculation: $(21.10 + 21.15) \div 2 = 21.125$ / 21.13 / 21.1 (cm ³) Do not award 21.10 Do not award 21.15	(1)

Question Number	Answer	Additional Guidance	Mark
3(c)	• moles calculated • concentration calculated	(1) (1) Example of a calculation: $5.00 \div 116 = 0.043103$ (mol) $0.043103 \div 0.25 = 0.17241$ / 0.17 (mol dm ⁻³) Ignore SF except 1SF TE M1 to M2	(2)

Question Number	Answer	Additional Guidance	Mark
3(d)	- moles of maleic acid (1) - use of ratio and moles of NaOH (1) - concentration of sodium hydroxide (1)	Example of a calculation: $(20 \div 1000) \times 0.17241 = 0.0034482 / 3.4482 \times 10^{-3} \text{ (mol)}$ 2:1 ratio so $0.0068964 / 6.8964 \times 10^{-3} \text{ mol NaOH}$ $0.0068964 \div (21.125 \div 1000) = 0.32646 / 0.33 \text{ mol dm}^{-3}$ TE throughout and on 3(b) and 3(c) Ignore SF except 1SF	(3)

Question Number	Answer	Additional Guidance	Mark
3(e)(i)	A description that makes reference to the following points: (shaken with) bromine water (1) - will decolourise with maleic acid and will not change colour with succinic acid (1)	Allow bromine Accept goes from brown/orange/yellow to colourless Allow red-brown to colourless Allow acidified potassium manganate(VII) for M1, will decolourise with maleic acid (M2) and no change with succinic acid	(2)

Question Number	Answer	Additional Guidance	Mark
3(e)(ii)	An answer that makes reference to the following point: • Z-but(-2-)enedioic acid	Allow <i>cis</i>-but(-2-)enedioic acid Allow Z-2-butene-1,4-dioic acid Allow <i>cis</i>-but-2-enedioic acid	(1)

(Total for Question 3 = 11 marks)

Question Number	Answer	Additional Guidance	Mark
4(a)	An answer that makes reference to the following points: 40-60°C - justification	(1) Allow any temperature in this range (1) The solution needs to be warm but if too hot presents a scald risk Allow so the organic liquid doesn't boil/evaporate Allow suggestions relating to rate needing to be increased Ignore responses based on time Marks are independent	(2)

Question Number	Answer	Additional Guidance	Mark
4(b)	An answer that makes reference to the following points: 1-chlorobutane and 1-bromobutane - all halogenoalkanes must be primary (to allow comparison) OR molecules must have the same chain length (to ensure similar solubility/miscibility/inductive effect)	(1) Accept CH₃CH₂CH₂CH₂Cl and CH₃CH₂CH₂CH₂Br Ignore C₄H₉Cl and C₄H₉Br (1) Allow same type/class/structure of halogenoalkane Allow only the halogen changes, not the alkane	(2)

Question Number	Answer	Additional Guidance	Mark
4(c)	An answer that makes reference to the following points: • $1 \div \text{time}$	Allow $1/\text{time}$ or reciprocal of time or one divided by time	(1)

Question Number	Answer	Additional Guidance	Mark
4(d)	An answer that makes reference to the following point: • the data (is discontinuous and) already shows the trend	Accept the percentage/experimental error/uncertainty in measuring the times is small in comparison with the differences between them. Allow the values are very different so the trend is clear Allow the three haloalkane rates are compared under identical conditions.	(1)


Question Number	Answer	Additional Guidance	Mark
4(e)	An answer that makes reference to the following point: • ammonia / NH_3	Allow NH_4OH Ignore references to concentration Do not award $\text{NH}_3 \cdot \text{H}_2\text{O}$	(1)

(Total for Question 4 = 7 marks)

Question Number	Answer	Additional Guidance	Mark
5(a)	An answer that makes reference to the following points: letters in the correct order	(2) A <u>F</u> <u>B</u> <u>D</u> <u>E</u> <u>C</u> All five in the correct order scores (2) Any two letters in the correct place (1)	(2)

Question Number	Answer	Additional Guidance	Mark
5(b)(i)	correct equation	Example of an equation: $\text{NaHCO}_3(\text{aq}) + \text{HCl}(\text{aq}) \rightarrow \text{NaCl}(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g})$ Allow ionic equation $\text{HCO}_3^-(\text{aq}) + \text{H}^+(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g})$	(2)
	correct state symbols	M2 dependent on M1 (or a near-miss)	(1)

Question Number	Answer	Additional Guidance	Mark
5(b)(ii)	An answer that makes reference to the following points: to dry the (organic) liquid	(1) Allow to absorb water Allow to act as a drying agent Allow dry the (organic) solution Allow “eliminate water” Ignore “remove water” alone Do not award dehydration	(1)

Question Number	Answer	Additional Guidance	Mark
5(c)(i)	An explanation that makes reference to the following points:  • alcohol X / /2-methylpropan-2-ol (1) • (alcohol X) is tertiary / 3° (1) • (alcohol X) will form a more stable carbocation (1)	If X, the formula and the name are given all must be correct Allow forms a tertiary carbocation Ignore shape of carbocation Ignore comments on activation energy	(3) Expert

Question Number	Answer	Additional Guidance	Mark
5(c)(ii)	A description that makes reference to the following points: • heat (with sulfuric acid and potassium dichromate(VI)) (1) • X will not turn green (1) • distil the products (from Y and Z) (1) • Z product will react with Fehling's/Benedict's (1) • to give a red/brown precipitate (1)	Allow warm Ignore reflux for M1 when aldehydes are implied Do not award hydrochloric acid Accept Y and Z will produce a colour change orange to green Allow X will not produce a colour change from orange Accept converse for Y product Accept Z product will react with Tollens' (1) to give a silver mirror (1) M5 has to have the correct reagent to be awarded "silver mirror" without Tollens would not score. Alternative: reflux (M1) M2 and M3 are the same as above Z product will react with $\text{CO}_3^{2-}/\text{HCO}_3^-$ (M4) To fizz / gas turns lime water milky/white (M5) Or Z product will react with PCl_5 (M4) To produce misty fumes (M5) Allow triiodomethane (iodoform) as a negative test for the product of Z (1), Y will give a pale-yellow precipitate (1)	(5) Expert

(Total for Question 5 = 13 marks)
TOTAL FOR PAPER = 50 MARKS