



Mark Scheme (Results)

Summer 2022

Pearson Edexcel International Advanced
Level in Chemistry (WCH15)
Paper 01: Transition Metals and Organic
Nitrogen Chemistry

Section A (multiple choice)

Question Number	Correct Answer	Mark
1(a)	The only correct answer is B (Fe^{2+}) <i>A is incorrect because Cl^- and $\text{S}_2\text{O}_8^{2-}$ ions are both negative so likely to repel</i> <i>C is incorrect because Cu^{2+} cannot oxidise I^-</i> <i>D is incorrect because Cu^+ cannot oxidise I^-</i>	(1)

Question Number	Correct Answer	Mark
1(b)	The only correct answer is D (homogeneous) <i>A is incorrect because the catalyst is not a product of the reaction</i> <i>B is incorrect because the catalyst is not an enzyme</i> <i>C is incorrect because the catalyst is not in a different physical state to the reactants</i>	(1)

Question Number	Correct Answer	Mark
1(c)	The only correct answer is D (sulfuric acid) <i>A is incorrect because ammonia is produced in industry using an iron catalyst</i> <i>B is incorrect because nitric acid is produced in industry using a platinum / rhodium catalyst</i> <i>C is incorrect because sodium hydroxide is produced in industry by electrolysis of brine, without a catalyst</i>	(1)

(Total for Question 1 = 3 marks)


Question Number	Correct Answer	Mark
2(a)	The only correct answer is A ($\text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^-$) <i>B is incorrect because it is the reverse of the reaction at the negative electrode</i> <i>C is incorrect because it is the reaction at the positive electrode</i> <i>D is incorrect because it is the reverse of the reaction at the positive electrode</i>	(1)

Question Number	Correct Answer	Mark
2(b)	The only correct answer is C (the catalyst is more efficient) <i>A is incorrect because the overall reaction is the same</i> <i>B is incorrect because the overall reaction is the same</i> <i>D is incorrect because the overall reaction is the same</i>	(1)

(Total for Question 2 = 2 marks)

Question Number	Correct Answer	Mark
3	The only correct answer is C (green) <i>A is incorrect because thiosulfate ions will reduce vanadate(V) to oxidation state = +3</i> <i>B is incorrect because thiosulfate ions will reduce vanadate(V) to oxidation state = +3</i> <i>D is incorrect because thiosulfate ions will not reduce vanadate(V) to oxidation state = +2</i>	(1)

(Total for Question 3 = 1 mark)

Question Number	Correct Answer	Mark
4	The only correct answer is C  <i>A is incorrect because it is not used in the treatment of cancer</i> <i>B is incorrect because it is not used in the treatment of cancer</i> <i>D is incorrect because it is the trans form of a complex used in the treatment of cancer</i>	(1)

(Total for Question 4 = 1 mark)

Question Number	Correct Answer	Mark
5	The only correct answer is A (NaOH(aq)) <i>B is incorrect because the $\text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O} \rightleftharpoons 2\text{CrO}_4^{2-} + 2\text{H}^+$ would move to the left on addition of acid</i> <i>C is incorrect because zinc would reduce $\text{Cr}_2\text{O}_7^{2-}$</i> <i>D is incorrect because hydrogen peroxide is used to oxidise Cr^{3+} to form $\text{Cr}_2\text{O}_7^{2-}$</i>	(1)

(Total for Question 5 = 1 mark)

Question Number	Correct Answer	Mark
6	The only correct answer is C (31.25) <i>A is incorrect because an incorrect expression to find uncertainty is used and only one burette reading is taken into account</i> <i>B is incorrect because only one burette reading is taken into account</i> <i>D is incorrect because it is simply the % uncertainty multiplied by 100</i>	(1)


Question Number	Correct Answer	Mark
7(a)	The only correct answer is B (62.5 %) <i>A is incorrect because the two carbonyl carbon atoms have not been included</i> <i>C is incorrect because the hydrogen atoms have not been included</i> <i>D is incorrect because two additional carbon atoms have been included</i>	(1)

(Total for Question 6 = 1 mark)

Question Number	Correct Answer	Mark
7(b)	The only correct answer is B (3) <i>A is incorrect because the carbons in the benzene ring are not all in the same environment</i> <i>C is incorrect because there are only 2 different carbon environments in the benzene ring</i> <i>D is incorrect because there are only 2 different carbon environments in the benzene ring</i>	(1)

Question Number	Correct Answer	Mark
7(c)	The only correct answer is A (a single type of monomer by an addition reaction) <i>B is incorrect because the polymer is not formed by a condensation reaction</i> <i>C is incorrect because the polymer is not formed by two different types of monomer</i> <i>D is incorrect because the polymer is not formed by two different types of monomer or a condensation reaction</i>	(1)

(Total for Question 7 = 3 marks)

Question Number	Correct Answer	Mark
8	The only correct answer is D  <i>A is incorrect because it is the phenylammonium ion</i> <i>B is incorrect because the bonding and charge is incorrect on the right hand nitrogen</i> <i>C is incorrect because the structure is an amine with a positive charge</i>	(1)

(Total for Question 8 = 1 mark)


Question Number	Correct Answer	Mark
9(a)	The only correct answer is B (ether) <i>A is incorrect because it would protonate the Grignard reagent</i> <i>C is incorrect because it is non-polar</i> <i>D is incorrect because it would react with the Grignard reagent to form a tertiary alcohol</i>	(1)

Question Number	Correct Answer	Mark
9(b)	The only correct answer is D (to ensure the solvent boils smoothly) <i>A is incorrect because the anti-bumping granules will not change the boiling temperature</i> <i>B is incorrect because this is the role of the condenser</i> <i>C is incorrect because the anti-bumping granules will not affect the flammability of the solvent</i>	(1)

Question Number	Correct Answer	Mark
9(c)	The only correct answer is D (negative and nucleophilic) <i>A is incorrect because the carbon atom is not positive or electrophilic</i> <i>B is incorrect because the carbon atom is not positive</i> <i>C is incorrect because the carbon atom is not electrophilic</i>	(1)

Question Number	Correct Answer	Mark
9(d)	The only correct answer is A (hexan-3-one) <i>B is incorrect because the product would be 2,4-dimethyloctan-4-ol</i> <i>C is incorrect because hexan-3-ol does not have a carbonyl bond</i> <i>D is incorrect because hexan-2-ol does not have a carbonyl bond</i>	(1)

(Total for Question 9 = 4 marks)

Question Number	Correct Answer	Mark
10(a)	The only correct answer is D  <i>A is incorrect because this ion would form in an acidic solution</i> <i>B is incorrect because this is the zwitterion</i> <i>C is incorrect because the OH group would not lose a proton</i>	(1)
10(b)	The only correct answer is B (ionic bonds) <i>A is incorrect because ionic bonds are far stronger than any hydrogen bonds</i> <i>C is incorrect because ionic bonds are far stronger than any London forces</i> <i>D is incorrect because the formation of a peptide bond forms a dipeptide</i>	(1)


(Total for Question 10 = 2 marks)

Question Number	Correct Answer	Mark
11	The only correct answer is A (more reactive and higher electron density) <i>B is incorrect because phenol has a higher electron density</i> <i>C is incorrect because phenol is not less reactive</i> <i>D is incorrect because phenol is not less reactive and has a higher electron density</i>	(1)

(Total for Question 11 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS

Section B

Question Number	Acceptable Answers	Additional Guidance	Mark
12(a)(i)	An answer that makes reference to the following points: - circle around arrow from $^+\text{CH}_3$ to ring (wrong direction) (1) - circle around arrow from bond attached to H, to partial ring (single-headed arrow) (1)	$\text{CH}_3\text{Cl} + \text{AlCl}_3 \longrightarrow ^+\text{CH}_3 + \text{AlCl}_4^-$  $\text{H}^+ + \text{AlCl}_4^- \longrightarrow \text{HCl} + \text{AlCl}_3$	(2)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(a)(ii)	An answer that makes reference to the following points: • arrow should move from ring / arrow is going the wrong way and ring is electron-rich / as ring cannot accept electrons / as $^+\text{CH}_3$ does not have a lone pair (of electrons) / as $^+\text{CH}_3$ needs to gain electrons / as $^+\text{CH}_3$ is an electrophile (1) • arrow (from C-H bond) should be double-headed and as both electrons in the bond pair move (to complete the ring) / as moving a single electron would not complete the ring / as moving a single electrons would form (free) radicals (1)	If no other credit awarded, then 1 mark can be given for both corrections correctly identified	(2)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(b)	An answer that makes reference to one of the following points: to prevent further substitutions (of nitro groups)	Allow methylbenzene is (more) reactive (than benzene) and because a methyl group is electron releasing Allow ring in methylbenzene is (more) electron-rich (than benzene) Allow forms dinitrobenzene / trinitrobenzene Ignore further reactions / forms other products	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(c)	An answer that makes reference to the following point: • oxidation	Allow oxidation and reduction / oxidation and redox Ignore references to redox	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(d)	An answer that makes reference to the following point: • Sn and (concentrated) HCl	Accept correct names Allow Zn as alternative to Sn Ignore references to concentration or state of HCl Ignore any references to heat / temperature Ignore addition of NaOH after reaction with Sn and HCl	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(e)	An explanation that makes reference to three of the following points: • (carbonyl) carbon is electron-deficient / has a partial positive charge • nitrogen (on NH₂ group) has a lone pair (of electrons) • which move to (carbonyl) carbon (to form bond) / which form a bond with the (carbonyl) carbon • Cl is a good leaving group / bond to Cl breaks / C-Cl bond pair moves to Cl	Allow M1, M2 and M3 on a clearly annotated diagram Ignore references to delocalisation Ignore references to nucleophilic substitution / nucleophilic addition-elimination	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
12(f)	- calculation of moles of 2-ethanoylaminobenzoic acid (1) - calculation of moles of benzene required, taking into account the overall percentage yield (1) - calculation of mass of benzene (1) - calculation of volume of benzene to 1/2/3 SF (1)	<u>Example of calculation</u> $5.92 \div 179 = 0.033073 \text{ (mol)} / 3.3073 \times 10^{-2}$ $0.033073 \times (100 \div 28.2) = 0.11728 \text{ (mol)}$ $0.11728 \times 78 = 9.1477 \text{ (g)}$ $9.1477 \div 0.879 = 10.407$ $= 10.4 / 10 \text{ cm}^3$ Ignore absence of units but do not award incorrect units. Marks 2 – 4 can be in any order Allow TE throughout Answer of 10 cm³ with no working scores M4 only Correct answer with some working scores 4	(4)


(Total for Question 12 = 14 marks)

Question Number	Acceptable Answers	Additional Guidance	Mark																				
13	This question assesses the student’s ability to show a coherent and logically structured answer with linkages and fully sustained reasoning. Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning. The following table shows how the marks should be awarded for indicative content. <table><tr><th>Number of indicative marking points seen in answer</th><th>Number of marks awarded for indicative marking points</th></tr><tr><td>6</td><td>4</td></tr><tr><td>5-4</td><td>3</td></tr><tr><td>3-2</td><td>2</td></tr><tr><td>1</td><td>1</td></tr><tr><td>0</td><td>0</td></tr></table> The following table shows how the marks should be awarded for structure and lines of reasoning <table><tr><th></th><th>Number of marks awarded for structure of answer and sustained lines of reasoning</th></tr><tr><td>Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout</td><td>2</td></tr><tr><td>Answer is partially structured with some linkages and lines of reasoning</td><td>1</td></tr><tr><td>Answer has no linkages between points and is unstructured</td><td>0</td></tr></table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure of answer and sustained lines of reasoning	Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2	Answer is partially structured with some linkages and lines of reasoning	1	Answer has no linkages between points and is unstructured	0	Guidance on how the mark scheme should be applied: The mark for indicative content should be added to the mark for lines of reasoning. For example, a response with four indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning). If there were no linkages between the points, then the same indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and zero marks for linkages). In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.	
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure of answer and sustained lines of reasoning																						
Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2																						
Answer is partially structured with some linkages and lines of reasoning	1																						
Answer has no linkages between points and is unstructured	0																						

	Indicative Points Similarities IP1 both alkalis initially react to give a green precipitate IP2 $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow [\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2] + 2\text{H}_2\text{O}$ / $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow [\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2] + 2\text{NH}_4^+$ IP3 these are deprotonation reactions Differences IP4 with excess ammonia (the green precipitate dissolves to) form a blue solution and no change with sodium hydroxide IP5 $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 6\text{NH}_3 \rightarrow [\text{Ni}(\text{NH}_3)_6]^{2+} + 6\text{H}_2\text{O}$ / $[\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2] + 6\text{NH}_3 \rightarrow [\text{Ni}(\text{NH}_3)_6]^{2+} + 4\text{H}_2\text{O}(\text{l}) + 2\text{OH}^-$ / $\text{Ni}(\text{OH})_2 + 6\text{NH}_3 \rightarrow \text{Ni}(\text{NH}_3)_6^{2+} + 2\text{OH}^-$ IP6 with excess ammonia it is ligand exchange Comment in equations allow use of round brackets instead of square brackets	Allow solid / crystals / ppt / ppte Allow $\text{Ni}^{2+} + 2\text{OH}^- \rightarrow \text{Ni}(\text{OH})_2$ $\text{NiSO}_4 + 2\text{NaOH} \rightarrow \text{Ni}(\text{OH})_2 + \text{Na}_2\text{SO}_4$ Allow acid-base reaction Ignore precipitation reaction / neutralisation reaction Do not award blue-green or blue-purple solution Allow $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 4\text{NH}_3 \rightarrow [\text{Ni}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 4\text{H}_2\text{O}$ Ignore state symbols in IP2 and IP5, even if incorrect Ignore omission of square brackets Allow ligand substitution	
--	---	---	--

(Total for Question 13 = 6 marks)

Question Number	Acceptable Answers	Additional Guidance	Mark																
14(a)	- calculation of mass of C and H (1) - calculation of mass of oxygen (1) - calculation of moles of C, H and O (1) - calculation of ratio and deduction of empirical formula (1)	<u>Example of calculation</u> <table border="1"> <thead> <tr> <th>Element</th><th>C</th><th>H</th><th>O</th></tr> </thead> <tbody> <tr> <td>Mass (g)</td><td>18.07 x (12÷44) = 4.9282</td><td>3.30 x (2÷18) = 0.36667</td><td>6.02 – (4.9282+0.36667) =0.72513</td></tr> <tr> <td>Moles (mol)</td><td>4.9282 ÷ 12 =0.41068</td><td>0.36667÷1 = 0.3667</td><td>0.72513 ÷ 16 = 0.045320</td></tr> <tr> <td>Ratio</td><td>0.41068÷0.045320 = 9.06</td><td>0.36667÷0.045320 = 8.09</td><td>0.045320÷0.045320 = 1</td></tr> </tbody> </table> C₉H₈O Allow TE throughout Ignore minor rounding errors in M1-M3	Element	C	H	O	Mass (g)	18.07 x (12÷44) = 4.9282	3.30 x (2÷18) = 0.36667	6.02 – (4.9282+0.36667) =0.72513	Moles (mol)	4.9282 ÷ 12 =0.41068	0.36667÷1 = 0.3667	0.72513 ÷ 16 = 0.045320	Ratio	0.41068÷0.045320 = 9.06	0.36667÷0.045320 = 8.09	0.045320÷0.045320 = 1	(4)
Element	C	H	O																
Mass (g)	18.07 x (12÷44) = 4.9282	3.30 x (2÷18) = 0.36667	6.02 – (4.9282+0.36667) =0.72513																
Moles (mol)	4.9282 ÷ 12 =0.41068	0.36667÷1 = 0.3667	0.72513 ÷ 16 = 0.045320																
Ratio	0.41068÷0.045320 = 9.06	0.36667÷0.045320 = 8.09	0.045320÷0.045320 = 1																

Question Number	Acceptable Answers	Additional Guidance	Mark
14(b)	An explanation that makes reference to the following points: • correct structure of Q shown (1) • sooty flame indicates benzene ring / arene / phenyl group / high C : H ratio / aromatic (1) • orange precipitate with 2,4-dinitrophenylhydrazine indicates carbonyl group / C=O / aldehyde or ketone (1) • silver precipitate with Tollens' reagent indicates aldehyde (1) • decolourises bromine water indicates C=C bond / alkene (functional group) / unsaturated (1) • exists as a pair of geometric isomers indicates only 1 hydrogen atom on each carbon of the C=C bond / each carbon of the C=C has two different groups attached (1)	e.g  Accept cis structure Accept skeletal structure without terminal hydrogen Accept hybrid structures e.g. partially displayed Ignore 'it is an alkene' Ignore references to phenol Do not award benzene Allow 'cannot have -CH=CH₂ group' Allow '2 different groups on each side of the C=C bond'	(6)

(Total for Question 14 = 10 marks)

Question Number	Acceptable Answers	Additional Guidance	Mark
15(a)(i)	$\text{Cr}_2\text{O}_7^{2-}$ orange and Cr^{3+} green	Do not award precipitate Do not award blue Ignore adjectives e.g. 'dark', 'pale' etc	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
15(a)(ii)	colour change (from orange to green at end-point) is not distinctive / colour change (from orange to green at end-point) is not sharp enough (without indicator) / colour change (at end-point) not easy to detect (without indicator)	Allow solutions (very) dilute so colour change hard to see (without indicator) Allow intense red-violet colour is not masked by colours of chromium species Allow idea that (all) chromium / iron species are coloured so the change is not easy to detect (without indicator)	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
15(a)(iii)	- calculation of moles of ammonium iron(II) sulfate (1) - calculation of moles of dichromate(VI) in titre (1) - calculation of moles of dichromate(VI) in original sample (1) - calculation of mass of potassium dichromate(VI) in original sample (1) - calculation of % by of potassium dichromate(VI) in 50 g of cement (1)	<u>Example of calculation</u> $3.24 \times 10^{-4} \times (10.90/1000) = 3.5316 \times 10^{-6} \text{ (mol)}$ $3.5316 \times 10^{-6} \div 6 = 5.8860 \times 10^{-7} \text{ (mol)}$ $5.8860 \times 10^{-7} \times 2 = 1.1772 \times 10^{-6} \text{ (mol)}$ $1.1772 \times 10^{-6} \times 294.2 = 3.4633 \times 10^{-4} \text{ (g)}$ $\frac{(3.4633 \times 10^{-4})}{50} \times 100 = 6.9266 \times 10^{-4} \%$ Allow TE throughout, but for M5 TE % must be less than 100 % Ignore SF Correct answer with or without working scores (5)	(5)

Question Number	Acceptable Answers	Additional Guidance	Mark
15(b)	An explanation that makes reference to the following points: - as it reacts with the COOH group to form COO⁻ / sodium carboxylate / a salt (1) - sodium salts are (more) soluble in water (than the acid) (1)	Allow ions are solvated by water / interact with water (more readily) Allow 'forms ionic substances which are more soluble in water'	(2)


Question Number	Acceptable Answers	Additional Guidance	Mark
15(c)(i)	An explanation that makes reference to the following points: - lone pair(s) of electrons on nitrogen (atoms) (1) - lone pair on one of the nitrogen (atoms) on the left of the C=O and lone pair on one of the nitrogen (atoms) on the right of the C=O (1) - Which form 2 dative (covalent) bonds (to the chromium ion) (1)	Allow lone pairs shown on diagram for M1 and M2 Ignore references to lone pairs on oxygen Do not award M2 if four lone pairs are referenced unless it's clear that only 2 of them, one from either side of the carbonyl carbon, form the bonds Allow dative (covalent) bonds shown on diagram	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
15(c)(ii)	chromium(VI) has an empty d subshell	Allow empty d orbitals (plural) Allow empty d orbital (singular) if clarified by correct electron configuration of ion Ignore idea that d orbitals do not split	(1)


Question Number	Acceptable Answers	Additional Guidance	Mark
15(d)	An explanation that makes reference to the following points: (reaction has) 5 particles on the left, but 7 on the right (1) - so there is a (pronounced) increase in entropy (of the system) / change in entropy (of the system) is positive / $\Delta S_{\text{(system)}}$ is positive (1)	Allow more particles on the right hand side (than on the left hand side) / increase in number of moles (of particles) Do not award use of molecules for particles Do not award incorrect numbers of particles Allow so there is a (pronounced) increase in disorder (of the system) Ignore there is a (pronounced) increase in total entropy M2 dependent on M1 or near miss e.g. use of 'molecules' in M1, increasing but incorrect number of particles in M1	(2)

(Total for Question 15 = 15 marks)

Question Number	Acceptable Answers	Additional Guidance	Mark
16	An answer that makes reference to the following points: - formation of 2-bromobutane using HBr (1) - use of ethanolic KCN / alcoholic KCN / KCN(eth) (1) - formation of 2-methylbutanenitrile (1) - using HCl(aq) (1) - formation of 2-methylbutanoic acid (1) - formation of ethyl 2-methylbutanoate using ethanol and sulfuric acid (1) OR - formation of 2-bromobutane using HBr (1) - using magnesium in (dry) ether (1) - formation of sec-butyl magnesium bromide (1) - using CO₂ and HCl(aq) (1) - formation of 2-methylbutanoic acid (1) - formation of ethyl 2-methylbutanoate using ethanol and sulfuric acid (1)	See below Allow HCl and 2-chlorobutane Must be in context of attempted reaction with a haloalkane Allow CN⁻ / Ignore HCN Accept any strong mineral acid Allow H⁺ Allow any strong mineral acid Allow H⁺ Do not award just 'acid' Must be in context of attempted reaction with a haloalkane Allow H⁺ / acid work up Do not award just 'acid' Accept any strong mineral acid Allow any strong mineral acid Allow H⁺ Do not award just 'acid'	(6)



OR



Note - the reaction of the Grignard with CO_2 and HCl can be shown as two separate steps. If this is the case ignore the structure of any intermediate species

(Total for Question 16 = 6 marks)

(TOTAL FOR SECTION B = 51 MARKS)

Question Number	Acceptable Answers	Additional Guidance	Mark
17(a)(i)	- calculation of moles of CuSCN (1) - calculation of moles of Cu (1) - calculation of mass of Cu (1) - calculation of % of Cu and statement that it is a gilding metal (1) Note – allow 121.5 for M_r of CuSCN in M1 but do not award 64 for copper in M3	<u>Example of calculation</u> $4.69 \div 121.6 = 0.038569 / 3.8569 \times 10^{-2} \text{ (mol)}$ 1:1 so = $0.038569 / 3.8569 \times 10^{-2} \text{ (mol)}$ M2 can be subsumed as part of M3 $0.038569 \times 63.5 = 2.4491 \text{ (g)}$ $(2.4491 \div 2.72) \times 100 = 90.04 \%$ Ignore SF except 1 SF Allow TE at each step Note – if TE for M4 gives answer outside range of 95-89 % then must be identified as NOT a gilding metal Do not award TE for M4 if answer > 100 % Allow calculation of % of Zn to show whether sample is a gilding metal	(4)


Question Number	Acceptable Answers	Additional Guidance	Mark
17(a)(ii)	An explanation that makes reference to the following points: - the E_{cell} data indicates that Cu^{2+} should not be reduced to Cu^+ and Cu^+ should be reduced to Cu (1) Cu^{2+} can be reduced to Cu^+ as the conditions must be non-standard and as the $E^\ominus$ values are so close (1) - but Cu^+ is not reduced to Cu as the reaction must be kinetically hindered / have a high activation energy / very slow (1)	Accept $E_{\text{cell}} = -0.02 \text{ V}$ and $E_{\text{cell}} = +0.35 \text{ V}$ Allow addition of OH^- ions as alternative for conditions must be non-standard Allow 'not kinetically favoured'	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
17(b)	- white precipitate forms / precipitate of $\text{Zn}(\text{H}_2\text{O})_4(\text{OH})_2$ forms (1) $\text{Zn}(\text{H}_2\text{O})_6^{2+} + 2\text{OH}^- \rightarrow \text{Zn}(\text{H}_2\text{O})_4(\text{OH})_2 + 2\text{H}_2\text{O}$ (1) - but as excess $\text{NaOH}(\text{aq})$ is added, precipitate will dissolve (1) - due to formation of $\text{Zn}(\text{OH})_4^{2-}$ (1)	Accept sufficient NaOH will need to be added to neutralise the excess nitric acid Allow precipitate of $\text{Zn}(\text{OH})_2$ forms / precipitate of zinc hydroxide forms M1 can be awarded from correct formulae and state symbol in equation, hence fully correct equation with state symbol for solid scores M1 and M2 Allow solid / crystals for ppt Allow $\text{Zn}^{2+} + 2\text{OH}^- \rightarrow \text{Zn}(\text{OH})_2$ Correct formulae for M4 can be shown as part of an equation, even if equation is not correct Ignore state symbols even if incorrect Ignore omission of square brackets Ignore comments on validity of procedure	(4)

Question Number	Acceptable Answers	Additional Guidance	Mark
17(c)	An explanation that makes reference to the following points: - zinc ions disrupt layers / disrupt structure of copper ions (1) (so) layers (of copper ions) are less likely to slide over each other (1)	Allow reference to atoms zinc ions are a different size to copper ions / zinc ions are larger than copper ions Do not award M1 if particles referred to as molecules or forces referred to as intermolecular forces Allow 'atoms are less likely to slide over each other' Allow reverse argument	(2)

Question Number	Acceptable Answers	Additional Guidance	Mark
17(d)(i)	A diagram that makes reference to the following points: - two correctly labelled electrodes (1) - both solutions and concentrations correct (1) - salt bridge labelled, touching both solutions and voltmeter shown (1)	Allow any soluble zinc and copper salts Allow name or formulae in M1 and M2 If the solution for the salt bridge is discussed it must be correct Ignore temperature and pressure	(3)

Example of diagram



Question Number	Acceptable Answers	Additional Guidance	Mark
17(d)(ii)	- calculation of E for zinc half cell / calculation of $E - E^\ominus$ (1) - insert values in Nernst equation and rearrangement of Nernst equation (1) - calculation of [ion] (1) Alternative Route - calculation of $E^\ominus_{\text{cell}}$ (1) - insert values in Nernst equation and rearrangement of Nernst equation (1) - calculation of [ion] (1)	<u>Example of calculation</u> $0.34 - 1.09 = -0.75 \text{ (V)}$ $-0.75 = -0.76 + (0.0260/2) \times \ln[\text{Zn}^{2+}]$ $\ln[\text{ion}] = (0.01) \times (2/0.026)$ $\ln [\text{ion}] = 0.769230$ $[\text{ion}] = 2.1581 \text{ (mol dm}^{-3}\text{)}$ Ignore SF except 1SF Allow TE throughout $0.34 - -0.76 = 1.10 \text{ (V)}$ $1.10 = 1.09 + (0.0260/2) \times \ln[\text{Zn}^{2+}]$ $\ln[\text{ion}] = (0.01) \times (2/0.026)$ $\ln [\text{ion}] = 0.769230$ $[\text{ion}] = 2.1581 \text{ (mol dm}^{-3}\text{)}$ Ignore SF except 1SF Allow TE throughout	(3)

(Total for Question 17 = 19 marks)
(TOTAL FOR SECTION C = 19 MARKS)
TOTAL FOR PAPER = 90 MARKS