



Mark Scheme (Results)

Summer 2023

Pearson Edexcel International Advanced
Subsidiary Level In Chemistry (WCH15)

Paper 01

Unit 5: Transition Metals and Organic Nitrogen
Chemistry

Section A

Question Number	Answer	Mark
1	The only correct answer is D $\text{Pt(s)} \text{V}^{2+}(\text{aq}), \text{V}^{3+}(\text{aq}) \text{Cu}^{2+}(\text{aq}) \text{Cu(s)}$ <i>A is not correct because the $\text{V}^{3+}(\text{aq})/\text{V}^{2+}(\text{aq})$ half-cell should have a platinum electrode and should show oxidation and the $\text{Cu}^{2+}(\text{aq})/\text{Cu(s)}$ half-cell should show reduction</i> <i>B is not correct because the $\text{V}^{3+}(\text{aq})/\text{V}^{2+}(\text{aq})$ half-cell should have a platinum electrode</i> <i>C is not correct because the $\text{V}^{3+}(\text{aq})/\text{V}^{2+}(\text{aq})$ half-cell should show oxidation and the $\text{Cu}^{2+}(\text{aq})/\text{Cu(s)}$ half-cell should show reduction</i>	(1) Computer

Question Number	Answer	Mark
2	The only correct answer is D $\text{Mg} + 2\text{Ce}^{4+} \rightarrow \text{Mg}^{2+} + 2\text{Ce}^{3+}$ <i>A is not correct because Ce is a weaker reducing agent than Mg</i> <i>B is not correct because Ce^{3+} is a weaker reducing agent than Ce</i> <i>C is not correct because Mn^{2+} is a weaker reducing agent than Mn</i>	(1) Computer

Question Number	Answer	Mark
3	The only correct answer is D $(\Delta S^{\circ}_{\text{total}})$ <i>A is not correct because E°_{cell} is directly proportional to $\ln K_c$</i> <i>B is not correct because E°_{cell} is directly proportional to $\Delta S^{\circ}_{\text{total}}$ and not ΔH°</i> <i>C is not correct because E°_{cell} is directly proportional to $\Delta S^{\circ}_{\text{total}}$ and not $\Delta S^{\circ}_{\text{system}}$</i>	(1) Computer

Question Number	Answer	Mark
4	The only correct answer is C (the reactants are thermodynamically unstable with respect to the products) <i>A is not correct because the reaction is thermodynamically feasible so will occur under certain conditions</i> <i>B is not correct because the $E^\ominus_{\text{cell}}$ value is a thermodynamic and not a kinetic property</i> <i>D is not correct because the reaction may be kinetically inert and the conditions may be non-standard</i>	(1) Computer

Question Number	Answer	Mark
5	The only correct answer is A ($\text{H}_2 + 2\text{OH}^- \rightarrow 2\text{H}_2\text{O} + 2\text{e}^-$) <i>B is not correct because hydrogen is consumed and not produced in a hydrogen-oxygen fuel cell</i> <i>C is not correct because oxygen is reduced at the positive electrode in a hydrogen-oxygen fuel cell</i> <i>D is not correct because oxygen is consumed and not produced in a hydrogen-oxygen fuel cell</i>	(1) Computer

Question Number	Answer	Mark
6	The only correct answer is D ($\text{Cu} [\text{Ar}]3\text{d}^{10}4\text{s}^1$) <i>A is not correct because the 4s electrons are removed before the 3d electrons</i> <i>B is not correct because the electronic configuration of chromium is $[\text{Ar}]3\text{d}^54\text{s}^1$</i> <i>C is not correct because the 4s electrons are removed before the 3d electrons</i>	(1) Computer

Question Number	Answer	Mark
7	The only correct answer is C (six) <i>A is not correct because it only takes into account water ligands</i> <i>B is not correct because it only takes into account ethanoate ions</i> <i>D is not correct because the coordination numbers of the two chromiums have been added together</i>	(1) Computer

Question Number	Answer	Mark
8	The only correct answer is A ($[\text{Fe}(\text{CN})_6]^{4-}$) <i>B is not correct because $\text{C}_2\text{O}_4^{2-}$ is a bidentate ligand</i> <i>C is not correct because EDTA^{4-} is a hexadentate ligand</i> <i>D is not correct because $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ is a bidentate ligand</i>	(1) Computer

Question Number	Answer	Mark
9	The only correct answer is A ($[\text{CuCl}_4]^{2-}$) <i>B is not correct because this complex is octahedral with a bond angle of 90°</i> <i>C is not correct because this complex is linear with a bond angle of 180°</i> <i>D is not correct because this complex is square planar with a bond angle of 90°</i>	(1) Computer

Question Number	Answer	Mark
10	The only correct answer is B ($[\text{Co}(\text{H}_2\text{O})_6]^{2+}$) <i>A is not correct because VO^{2+} is blue</i> <i>C is not correct because $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$ is blue</i> <i>D is not correct because $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ is blue</i>	(1) Computer

Question Number	Answer	Mark
11	The only correct answer is D ($2\text{Cr}(\text{OH})_3 + 3\text{H}_2\text{O}_2 + 4\text{KOH} \rightarrow 2\text{K}_2\text{CrO}_4 + 8\text{H}_2\text{O}$) <i>A is not correct because FeCl_2 forms a green solution</i> <i>B is not correct because this is not a redox reaction</i> <i>C is not correct because this is not a redox reaction</i>	(1) Computer

Question Number	Answer	Mark
12	The only correct answer is C ($[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 4\text{H}_2\text{O}$) <i>A is not correct because this is the ionic equation describing the deprotonation when ammonia is not in excess</i> <i>B is not correct because four water ligands are exchanged by ammine ligands when ammonia is in excess</i> <i>D is not correct because four water ligands are exchanged by ammine ligands when ammonia is in excess</i>	(1) Computer


Question Number	Answer	Mark
13	The only correct answer is A ($[\text{Zn}(\text{OH})_4]^{2-} + 2\text{H}_3\text{O}^+ \rightarrow [\text{Zn}(\text{H}_2\text{O})_4(\text{OH})_2]$) <i>B is not correct because $[\text{Zn}(\text{OH})_4]^{2-}$ is a soluble complex ion</i> <i>C is not correct because $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ is a soluble complex ion</i> <i>D is not correct because $[\text{Cr}(\text{OH})_6]^{3-}$ is a soluble complex ion</i>	(1) Computer


Question Number	Answer	Mark
14	The only correct answer is C (+5 $\rightarrow$ +4 $\rightarrow$ +5) <i>A is not correct because the oxidation state in V_2O_5 is +5 not +2</i> <i>B is not correct because the oxidation state in V_2O_5 is +5 not +2</i> <i>D is not correct because the vanadium cannot be oxidised from +5 to +6</i>	(1) Computer


Question Number	Answer	Mark
15	The only correct answer is C (14) <i>A is not correct because each carbon contributes one electron from a p-orbital</i> <i>B is not correct because each carbon contributes one electron from a p-orbital</i> <i>D is not correct because each carbon contributes one electron from a p-orbital</i>	(1) Computer

Question Number	Answer	Mark
16	The only correct answer is C (a lone pair of electrons on oxygen in phenol is delocalised into the ring) <i>A is not correct because the polarity of the O-H bond does not increase the electron density of the benzene ring</i> <i>B is not correct because the electronegativity of the oxygen atom does not increase the electron density of the benzene ring</i> <i>D is not correct because there is a greater electron density in the ring in phenol than in benzene</i>	(1) Computer

Question Number	Answer	Mark
17	The only correct answer is B ($\text{C}_2\text{H}_5\text{NH}_2 \xrightarrow{\text{C}_2\text{H}_5\text{Br}}$) <i>A is not correct because the reduction of a nitrile forms a primary amine</i> <i>C is not correct because the products would be a tertiary amine and a quaternary ammonium salt</i> <i>D is not correct because the alkaline hydrolysis of this amide forms a primary amine</i>	(1) Computer

Question Number	Answer	Mark
18	The only correct answer is D () <i>A is not correct because this is not an azo dye</i> <i>B is not correct because this is not an azo dye</i> <i>C is not correct because this azo dye could only form if the reagents were not in excess</i>	(1) Computer

Question Number	Answer	Mark
19	The only correct answer is B () <i>A is not correct because this amino acid contains one acidic group and one basic group</i> <i>C is not correct because this amino acid contains one acidic group and two basic groups</i> <i>D is not correct because this amino acid contains one acidic group and two basic groups</i>	(1) Computer

Question Number	Answer	Mark
20	The only correct answer is A () <i>B is not correct because these reagents would lead to the formation of compound B</i> <i>C is not correct because these reagents would lead to the formation of compound B</i> <i>D is not correct because these reagents would lead to the formation of compound B</i>	(1) Computer

Total for Section A = 20 marks

Section B

Question Number	Answer	Additional Guidance	Mark
21(a)	An explanation that makes reference to the following points: • (Hg^+ is $[\text{Xe}]4f^{14} 5d^{10}6s^1$ and (Hg^{2+} is $[\text{Xe}]4f^{14} 5d^{10}(6s^0)$ (1) • (d-block element as last) electron goes into a (5)d-orbital(s) (when the electronic configuration is written according to the Aufbau principle) (1) • (not transition element as) Hg^+ and Hg^{2+}/(stable) ions do not have incompletely filled (5)d-orbital(s) (1)	Accept use of d-subshell for d-orbital(s) Allow use of d-shell for d-subshell Penalise use of just d-block for d-shell once only Penalise use of 3d/4d for 5d once only Allow Hg loses (only) its 6s electrons (when forming ions/compounds) Do not award answer in terms of the electronic configuration of an ion of mercury Allow Hg^+ and Hg^{2+}/(stable) ions have completely full (5)d-orbital(s) Ignore any reference to d-d transitions / other transition element properties Do not award answer in terms of the electronic configuration of the element / an Hg atom	(3) Expert

Question Number	Answer	Additional Guidance	Mark
21(b)(i)	An explanation that makes reference to the following points: - Hg/mercury oxidised and from 0 (in Hg) to +2 (in $\text{Hg}(\text{NO}_3)_2$) (1) - N/nitrogen is reduced and from +5 (in HNO_3) to +2 (in NO) (1)	Allow oxidation numbers from annotation to the equation Ignore any reference to electron loss/gain Do not award reference to oxidation of any other element Do not award HNO_3 is reduced Do not award reference to reduction of any other element If no other mark awarded, Hg/mercury oxidised and N/nitrogen reduced OR Hg/mercury from 0 to +2 and N/nitrogen from +5 to +2 scores (1)	(2) Expert

Question Number	Answer	Additional Guidance	Mark
21(b)(ii)	An answer that makes reference to the following points: - ionic half-equation for oxidation of mercury (1) - ionic half-equation for reduction of nitrate (1)	Allow multiples and $\rightleftharpoons$ for $\rightarrow$ Ignore state symbols, even if incorrect Examples of ionic half-equations: $\text{Hg} \rightarrow \text{Hg}^{2+} + 2\text{e}^{(-)}$ Allow $\text{Hg} - 2\text{e}^{(-)} \rightarrow \text{Hg}^{2+}$ Do not award half-equation including $\text{HNO}_3/\text{NO}_3^-$ $4\text{H}^+ + \text{NO}_3^- + 3\text{e}^{(-)} \rightarrow \text{NO} + 2\text{H}_2\text{O}$ Allow $3\text{H}^+ + \text{HNO}_3 + 3\text{e}^{(-)} \rightarrow \text{NO} + 2\text{H}_2\text{O}$ Allow $4\text{HNO}_3 + 3\text{e}^{(-)} \rightarrow \text{NO} + 2\text{H}_2\text{O} + 3\text{NO}_3^-$	(2) Expert

Question Number	Answer	Additional Guidance	Mark
21(b)(iii)	An answer that makes reference to the following point: balanced equation	Example of completed equation: $3\text{Hg(l)} + 8\text{HNO}_3\text{(aq)} \rightarrow 3\text{Hg(NO}_3)_2\text{(aq)} + 2\text{NO(g)} + 4\text{H}_2\text{O(l)}$ Allow multiples	(1) Clerical


Question Number	Answer	Additional Guidance	Mark
21(c)(i)	An answer that makes reference to the following point: • correct species • balanced equation	Example of completed equation: $\text{Hg}(\text{NO}_3)_2 + 3\text{C}_2\text{H}_5\text{OH} \rightarrow \text{Hg}(\text{CNO})_2 + 2\text{CH}_3\text{CHO} + 5\text{H}_2\text{O}$ Ignore state symbols even if incorrect (1) Do not award molecular formulae eg $\text{C}_2\text{H}_4\text{O}$ for CH_3CHO Do not award CH_3COH for CH_3CHO (1) Allow multiples No TE on M1 except on correct molecular formulae and on CH_3COH	(2) Graduate

Question Number	Answer	Additional Guidance	Mark
21(c)(ii)	• moles of $\text{Hg}(\text{CNO})_2$ • moles of gas produced • volume of gas produced	Correct answer with some working scores (3) Ignore SF except 1SF throughout Example of calculation: (1) $n = \frac{1.00}{284.6} = 0.0035137 / 3.5137 \times 10^{-3}$ (1) $n = 0.0035137 \times 2 = 0.0070274 / 7.0274 \times 10^{-3}$ TE on M1 (1) $v = 0.0070274 \times 24000 = 168.66 \text{ (cm}^3\text{)}$ Accept 0.16866 dm^3 TE on M2	(3) Expert

Question Number	Answer	Additional Guidance	Mark
21(d)(i)	An answer that makes reference to the following point: (to provide a) constant concentration (of Cl^-)	Allow KCl for Cl^- throughout Allow to keep the solution / Cl^- saturated Allow to replace Cl^- Ignore just to provide Cl^- Ignore stated concentrations Do not award salt bridge / to complete the circuit Do not award catalyst	(1) Expert

Question Number	Answer	Additional Guidance	Mark
21(d)(ii)	An answer that makes reference to the following point: $(0.24 - 0.37 =) -0.13 \text{ (V)}$	Ignore working, even if incorrect	(1) Expert

Question Number	Answer	Additional Guidance	Mark
21(d)(iii)	An answer that makes reference to the following point: $\text{Hg}_2\text{Cl}_2 + \text{Sn} \rightarrow 2\text{Hg} + \text{Sn}^{2+} + 2\text{Cl}^-$	Allow $\text{Hg}_2\text{Cl}_2 + \text{Sn} \rightarrow 2\text{Hg} + \text{SnCl}_2$ Allow multiples Allow $\rightleftharpoons$ for $\rightarrow$ Ignore state symbols even if incorrect Ignore half-equations even if incorrect Ignore use of cell diagrams Do not award uncanceled electrons Do not award $2\text{Hg}^+ (+ 2\text{Cl}^-)$ for Hg_2Cl_2 If answer to (d)(ii) is +0.61 (V) / +0.37 (V) / greater than +0.24 (V), equation must be reversed: $2\text{Hg} + \text{Sn}^{2+} + 2\text{Cl}^- \rightarrow \text{Hg}_2\text{Cl}_2 + \text{Sn}$ OR $2\text{Hg} + \text{SnCl}_2 \rightarrow \text{Hg}_2\text{Cl}_2 + \text{Sn}$	(1) Expert

Question Number	Answer	Additional Guidance	Mark
21(d)(iv)	An answer that makes reference to the following points: - hydrogen (gas)/H₂((g)) and 100 kPa - platinum (solid)/Pt((s)) and 298 K / 25°C - hydrochloric acid/HCl((aq)) and 1 mol dm⁻³	Example of completed diagram:  (1) Allow 101 kPa / 100 000 Pa / 101 000 Pa / 1 atm / 1 bar Accept platinum black for platinum Ignore porous (1) Ignore omission of state symbol 298 K / 25°C may be shown anywhere Allow H⁺ / H₃O⁺ for hydrochloric acid (1) Allow M for mol dm⁻³ Do not award 0.5 mol dm⁻³ H₂SO₄ If no other mark awarded: hydrogen/H₂ and hydrochloric acid/HCl/H⁺/H₃O⁺ and platinum/Pt scores (1)	(3) Graduate

Question Number	Answer	Additional Guidance	Mark
21(d)(v)	An answer that makes reference to any one of the following points: • (calomel electrode) does not require a (separate) salt bridge OR (calomel electrode) does not require a continuous supply of hydrogen / gas OR platinum/Pt (of hydrogen electrode is) easily poisoned OR difficult to ensure hydrogen electrode is at equilibrium	Ignore calomel electrode is quicker to use / easier to set up / done in the same container / more portable Accept does not require a hydrogen / gas generator Ignore just does not require hydrogen / gas Ignore any reference to pressure Ignore hydrogen is flammable / explosive / difficult to store Ignore (calomel electrode is) safer Ignore platinum is expensive Ignore (calomel electrode) is cheaper Allow (calomel electrode) reaches equilibrium sooner Allow (calomel electrode gives) more stable (reading) Ignore (calomel electrode is) more accurate Ignore calomel electrode potential is more positive	(1) Expert

(Total for Question 21 = 20 marks)

Question Number	Answer	Additional Guidance	Mark
22	An answer that makes reference to the following points: - moles of FeSO_4 - moles of excess MnO_4^- - initial moles of MnO_4^- and moles of MnO_4^- reacted - moles of $\text{C}_2\text{O}_4^{2-}$ - mass of CaC_2O_4 - percentage by mass of CaC_2O_4 and answer to 2SF or 3SF	Correct answer to 2SF or 3SF with some working scores (6) Ignore SF except 1SF Example of calculation: (1) $n = 0.0500 \times \frac{25.95}{1000} = 0.0012975 / 1.2975 \times 10^{-3}$ (1) $n = 0.0012975 \div 5 = 0.0002595 / 2.595 \times 10^{-4}$ TE on moles of FeSO_4 $n = 0.0100 \times \frac{50.0}{1000} = 0.0005 / 5 \times 10^{-4}$ (Allow 1 SF) and (1) $n = 0.0005 - 0.0002595 = 0.0002405 / 2.405 \times 10^{-4}$ TE on moles of excess MnO_4^- provided answer is positive (1) $n = 0.0002405 \times 2.5 = 0.00060125 / 6.0125 \times 10^{-4}$ TE on moles of MnO_4^- reacted (1) mass = $0.00060125 \times 128.1 = 0.077020$ (g) TE on moles of $\text{C}_2\text{O}_4^{2-}$ (1) % mass = $\frac{0.077020}{11.4} \times 100 = 0.67562$ (%) = 0.68 / 0.676 (%) TE on mass of CaC_2O_4 provided positive value to 2SF/3SF and < 100% Allow use of 128 for M_r of CaC_2O_4 giving 0.675 (%)	(6) Expert


(Total for Question 22 = 6 marks)

Question Number	Answer	Additional Guidance	Mark																				
*23	This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning. Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning. The following table shows how the marks should be awarded for indicative content. <table><tr><th>Number of indicative marking points seen in answer</th><th>Number of marks awarded for indicative marking points</th></tr><tr><td>6</td><td>4</td></tr><tr><td>5-4</td><td>3</td></tr><tr><td>3-2</td><td>2</td></tr><tr><td>1</td><td>1</td></tr><tr><td>0</td><td>0</td></tr></table> The following table shows how the marks should be awarded for structure and lines of reasoning. <table><tr><th></th><th>Number of marks awarded for structure and sustained lines of reasoning</th></tr><tr><td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td><td>2</td></tr><tr><td>Answer is partially structured with some linkages and lines of reasoning.</td><td>1</td></tr><tr><td>Answer has no linkages between points and is unstructured.</td><td>0</td></tr></table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and sustained lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning). If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages). If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded, do not deduct mark(s). Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.	(6) Expert
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure and sustained lines of reasoning																						
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2																						
Answer is partially structured with some linkages and lines of reasoning.	1																						
Answer has no linkages between points and is unstructured.	0																						



	Indicative points: • IP1: thermochemical data calculation (enthalpy of hydrogenation of 1,3,5-cyclohexatriene / benzene is) expected to be $-360 \text{ (kJ mol}^{-1}\text{)}$ • IP2: thermochemical data comparison (enthalpy of hydrogenation is) less exothermic / less negative than expected (for 1,3,5-cyclohexatriene) less exothermic / more stable by $152 \text{ (kJ mol}^{-1}\text{)}$ scores IP1 and IP2 • IP3: X-ray diffraction data (carbon-carbon) bond lengths in benzene are equal Ignore any reference to bond strength / bond angle in IP3 and IP4 • IP4: X-ray diffraction data (carbon-carbon) bond length in benzene is longer than (localised) C=C (in cyclohexene) • IP5: Bromination data (product for benzene is formed by electrophilic) substitution • IP6: Bromination data (benzene π-bonds less reactive than localised π-bonds and) requires (FeBr₃) catalyst (and heat)	Allow (enthalpy of hydrogenation is) expected to be three times the value for cyclohexene Allow (enthalpy of hydrogenation is) different by $152 \text{ (kJ mol}^{-1}\text{)}$ Accept reverse argument Ignore higher/lower for less exothermic Ignore benzene more exothermic than cyclohexene Ignore just benzene more stable than expected Do not award enthalpy required/needed Allow (carbon-carbon) bond lengths are not different Allow cyclohexene (carbon-carbon) bond lengths are different Accept (carbon-carbon) bond length is in between C=C and C–C (in cyclohexene) Allow (carbon-carbon) bond length is shorter than C–C (in cyclohexene) Allow (benzene) does not react by addition Allow cyclohexene/localised π-bonds react by addition Ignore any equations / mechanisms Do not award nucleophilic (substitution / addition) Accept cyclohexene does not require a catalyst Allow halogen carrier for catalyst Ignore just benzene does not decolourise bromine water Do not award Fe catalyst	
--	---	--	--

(Total for Question 23 = 6 marks)

Question Number	Answer	Additional Guidance	Mark
24(a)(i)	An answer that makes reference to the following point: prop-2-enamide / 2-propenamide	Allow capital letters and spaces Ignore omission of hyphen Allow propenamide Allow 'ene' for 'en' Allow propyl for prop Do not award propan for prop Do not award N- prefix Do not award cis/trans/E/Z- prefix	(1) Graduate

Question Number	Answer	Additional Guidance	Mark
24(a)(ii)	An explanation that makes reference to the following points: (PAM can form many) hydrogen bonds with water - H-bonds (with water) can form at NH_2 and $(\text{C}=\text{O})$ - diagram of (at least one) hydrogen bond between a water molecule and any amide group	(1) Allow M1 from a labelled diagram Ignore PAM reacts with water / acts as a base / accepts a proton from water / forms RNH_3^+ (1) M2 can be awarded from a diagram (1) diagram must include: H-bond to lone pair on O or N and $\delta^+\text{H}$ atom Ignore bond angle Do not award H-bond shown as coordinate bond / solid line (ie covalent bond) Example of diagram scoring (3):  Allow H-bond between lone pair on N of NH_2 and $\delta^+\text{H}$ of water	(3) Expert




Question Number	Answer	Additional Guidance	Mark
24(a)(iii)	An explanation that makes reference to the following points: - carboxylate / COO^- (above pH 8) (1) - repulsion between negative charges (above pH 8) (1)	Allow carboxylic acid/COOH/OH groups are deprotonated / donate H^+ / become anions Allow OH^- removes H atoms involved in hydrogen bonds Ignore just PAA is deprotonated / donates H^+ / becomes anion Ignore just salt is formed Do not award zwitterion is formed Allow (COO^-) cannot form (intramolecular) hydrogen bonds Allow (all) hydrogen bonds break Ignore hydrogen bonds weaken Ignore fewer hydrogen bonds Ignore any reference to denaturation Ignore any reference to intermolecular hydrogen bonds	(2) Expert



Question Number	Answer	Additional Guidance	Mark
24(b)(i)	An answer that makes reference to the following point: structure of vinylpyrrolidone monomer	Allow any combination of skeletal, structural or displayed formulae Examples of structure:  	(1) Expert

Question Number	Answer	Additional Guidance	Mark
24(b)(ii)	An answer that makes reference to the following points: - molar mass of vinylpyrrolidone monomer / PVP repeat unit - number of monomers per polymer and answer to nearest whole number	Correct answer with some working scores (2) Example of calculation: $M(C_6H_9NO) = 111.0 / 111 \text{ (g mol}^{-1}\text{)}$ TE on (b)(i) if molar mass is not 111.0 / 111 number = $90000 \div 111.0 = 810.81 = 811$ TE on M1	(2) Expert

Question Number	Answer	Additional Guidance	Mark
24(b)(iii)	An answer that makes reference to the following points: - mass of PVP per tablet - number of moles of PVP polymer - number of molecules of PVP polymer	Correct answer with some working scores (3) Example of calculation: (1) $\text{mass} = \frac{0.740}{100} \times 4.0 = 0.0296 \text{ (g)}$ Accept 29.6 (mg) Ignore SF except 1 SF (1) $\text{moles} = 0.0296 \div 90000 = 3.2889 \times 10^{-7}$ Allow $3.2889 \times 10^{-4} \text{ mmol}$ TE on M1 Ignore SF except 1 SF (1) $\text{molecules} = 3.2889 \times 10^{-7} \times 6.02 \times 10^{23} = 1.9799 \times 10^{17}$ Accept $2.0 \times 10^{17} / 2 \times 10^{17}$ TE on M2 (from any M_r value) Ignore SF Do not award multiplication of N_A by mass	(3) Expert

Question Number	Answer	Additional Guidance	Mark
24(c)(i)	An answer that makes reference to the following points: (polymer is a very) large molecule OR (polymer is formed from) large number of / many monomers (1) (condensation as) splitting off of a (small) molecule (1)	Allow long-chain (molecule) Allow macromolecule Allow repeating for many Ignore 2 or more / several / different for many Ignore (formed by) addition Allow with loss/elimination of H₂O/HCl Ignore forms byproduct	(2) Expert

Question Number	Answer	Additional Guidance	Mark
24(c)(ii)	An answer that makes reference to the following points: • structure of hexanedioyl dichloride • structure of hexane-1,6-diamine	Accept monomers in either order Allow any combination of skeletal, structural or displayed formulae If more than one type of formula shown, all must be correct Penalise errors in chain length once only Ignore connectivity in structural formulae eg Allow $\text{COClCH}_2\ldots$ / $\text{COOHCH}_2\ldots$ / $\text{NH}_2\text{CH}_2\ldots$ eg Allow $\text{C=OCl-CH}_2\ldots$ / $\text{C=OOH-CH}_2\ldots$ / $\text{NH}_2\text{-CH}_2\ldots$ Penalise connectivity in skeletal / displayed formulae once only Allow structure of hexanedioic acid Examples of correct structures: $\text{ClOCCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{COCl}$ / $\text{ClOC(CH}_2)_4\text{COCl}$  $\text{HOOCCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$ / $\text{HOOC(CH}_2)_4\text{COOH}$  $\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ / $\text{H}_2\text{N(CH}_2)_6\text{NH}_2$ 	(2) Expert

Question Number	Answer	Additional Guidance	Mark
24(c)(iii)	An answer that makes reference to the following points: • amide linkage (within polymer chain) (1) • two repeat units (1)	Allow any combination of skeletal, structural or displayed formulae Allow --NHCO-- / --CONH-- / --HNCO-- / --OCNH-- Ignore omission of square brackets Ignore n Examples of two repeat units:  $\text{--CO(CH}_2)_5\text{NHCO(CH}_2)_5\text{NH--}$  $\text{--NH(CH}_2)_5\text{CONH(CH}_2)_5\text{CO--}$	(2) Expert

(Total for Question 24 = 18 marks)


Total for Section B = 50 marks




Section C



Question Number	Answer	Additional Guidance	Mark
25(a)	An answer that makes reference to the following points: • ester (1) • (primary) amine and arene/benzene/phenyl OR phenylamine (1)	Ignore any structures / formulae Ignore carbonyl Do not award ketone / aldehyde / carboxylic acid Do not award ether Allow amino Allow aryl Ignore alkyl/alkane Do not award alkene Do not award phenol Allow aniline Allow aromatic amine	(2) Graduate

Question Number	Answer	Additional Guidance	Mark
25(b)(i)	An answer that makes reference to the following points: - molecular formulae of procaine and HCl (1) - molecular formula of procaine monohydrochloride (1)	Ignore non-molecular formulae $C_{13}H_{20}N_2O_2 + HCl$ Allow elements in any order $C_{13}H_{21}Cl^{(-)}N_2^{(+)}O_2$ Allow elements in any order TE on molecular formula of procaine Ignore position of charges Do not award separate $C_{13}H_{21}N_2^+O_2$ and Cl^- ions Do not award any additional product(s) Example of equation: $C_{13}H_{20}N_2O_2 + HCl \rightarrow C_{13}H_{21}ClN_2O_2$ scores (2)	(2) Graduate






Question Number	Answer	Additional Guidance	Mark
25(b)(ii)	An explanation that makes reference to the following points: - identification of tertiary amine nitrogen and effect of ethyl groups OR benzene ring EITHER ethyl / alkyl (groups) are electron donating OR lone pair (on N of NH₂ partially) delocalised into (aromatic) π-bond(s) (1) - second effect (1)	Ignore just comparison of electron density on N atoms Ignore just comparison of ability of (N) lone pairs to accept H⁺ Accept any unambiguous identification Accept ethyl / alkyl has positive inductive effect Allow ethyl / alkyl are electron pushing / electron releasing Allow methyl / R / attached groups for ethyl / alkyl Accept non-bonding pair for lone pair Allow electron pair for lone pair Allow overlaps with / interacts with / released into / drawn into for delocalised into Allow p-orbitals / ring for (aromatic) π-bond(s) Ignore just benzene for (aromatic) π-bond(s) Ignore just ring is electron withdrawing (with no mention of electron pair) If no other mark awarded, tertiary / aliphatic amine is more basic OR aromatic / primary amine is less basic scores (1)	(2) Expert

Question Number	Answer	Additional Guidance	Mark
25(c)	An answer that makes reference to the following point: • structure of 4-aminobenzoic acid	Allow any combination of skeletal, structural or displayed formulae Example of structure:  Allow Kekulé benzene Allow protonation of -NH_2 to -NH_3^+ Allow deprotonation of -COOH to -COO^- Allow zwitterion Do not award any other hydrolysis product	(1) Expert

Question Number	Answer	Additional Guidance	Mark
25(d)(i)	An answer that makes reference to the following points: - equation for formation of nitronium ion (1) - curly arrow from within hexagon to anywhere on NO_2^+ (1) - structure of intermediate ion (1) - curly arrow from C–H bond to within ring and correct product and H^+ (1)	Ignore omission or incorrect placement of methyl groups in M2 and M3 $\text{HNO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{NO}_2^+ + \text{HSO}_4^- + \text{H}_2\text{O}$ OR $\text{HNO}_3 + 2\text{H}_2\text{SO}_4 \rightarrow \text{NO}_2^+ + 2\text{HSO}_4^- + \text{H}_3\text{O}^+$ OR $\text{HNO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{NO}_3^+ + \text{HSO}_4^-$ and $\text{H}_2\text{NO}_3^+ \rightarrow \text{NO}_2^+ + \text{H}_2\text{O}$  TE on electrophile from M1 provided positively charged Do not award lone pair on N of NO_2^+  Allow any part of gap in 'horseshoe' facing tetrahedral carbon and covering at least three carbons with some part of positive sign within 'horseshoe'. 'Horseshoe' may be dashed TE on electrophile from M2 Do not award $\text{NO}_2\text{--C}$ connectivity Do not award dashed C–H and C–N bonds unless 3D structure 	(4) Expert

Question Number	Answer	Additional Guidance	Mark
25(d)(ii)	An answer that makes reference to the following points: • structure of 2,4-dimethylnitrobenzene (1) • structure of 3,5-dimethylnitrobenzene (1)	Allow structures with di-, tri- or tetranitro substitution Ignore connectivity of NO₂ group Penalise omission of delocalised ring once only Examples of structures:  	(2) Graduate

Question Number	Answer	Additional Guidance	Mark
25(d)(iii)	An answer that makes reference to the following point: • tin and (concentrated) hydrochloric acid	Accept Sn and HCl((aq)) Ignore heat / reflux Ignore NaOH in second step Do not award NaOH with Sn and HCl in the same step Do not award any reference to catalyst	(1) Graduate

Question Number	Answer	Additional Guidance	Mark
25(d)(iv)	An answer that makes reference to the following point: skeletal formula of 2-chloroethanoyl chloride	Ignore non-skeletal formulae Ignore bond angles and bond lengths  Allow skeletal formula of 2-chloroethanoic acid  Ignore OH connectivity Allow skeletal formula of 2-chloroethanoic anhydride  Allow skeletal formula of 2-chloroethanoyl bromide  Do not award skeletal formula of 2-chloroethanoyl cation 	(1) Graduate

Question Number	Answer	Additional Guidance	Mark
25(d)(v)	An answer that makes reference to the following point: nucleophilic substitution	Allow S_N2 / S_N1 Do not award any other answer	(1) Clerical

Question Number	Answer	Additional Guidance	Mark
25(e)	An answer that makes reference to the following point: indication of chiral centre	articaine Allow any indication Do not award any other answer	(1) Graduate

Question Number	Answer	Additional Guidance	Mark
25(f)	An answer that makes reference to the following points: - number of half-lives in 4 hours (1) - mass of articaine remaining in mg (1) - conversion of mg to μg (1)	Example of calculation: half-lives = $\frac{(4 \times 60)}{20} = 12$ mass = $100 \times 0.5^{12} = 0.024414$ (mg) TE on M1 Ignore SF except 1SF mass = $0.024414 \times 1000 = 24.414$ (μg) TE on M1 and M2	(3) Expert

(Total for Question 25 = 20 marks)

Total for Section C = 20 marks

Total for Paper = 90 marks