

Mark Scheme (Results)

January 2021

Pearson Edexcel International Advanced Level In Chemistry (WCH16) Paper 1 Practical Skills in Chemistry II

Question	Answer	Additional Guidance	Mark
number			
1(a)(i)	An answer that makes reference to the following points: • corrosive	Allow corrosive and acidic	(1)
		Do not award flammable, harmful, irritant, oxidising, oxidant, toxic	

Question number	Answer		Additional Guidance	Mark
1(a)(ii)	An answer that makes reference to the following points:		Use of a beaker / conical flask to add A will score (0)	(2)
	drop by drop	(1)	M1 dependent on apparatus which produces drops e.g burette, syringe	
			Allow dropwise / add in drops	
			Do not award add in small amounts	
	• (dropping) pipette / teat pipette	(1)	Allow burette	
			Do no award pipette with a stated volume such as 5 cm ³ Do not award just 'dropper'	

Question	Answer	Additional Guidance	Mark
number			
1(b)(i)	An answer that makes reference to the following point: • $Cu_{((aq))}^{2+} / Cu(H_2O)_6^{2+}$	Allow Copper(II) ions / hove gue copper(II)	(1)
		Allow Copper(II) ions / hexaquacopper(II) Do not award Cr ²⁺ / Cr ³⁺ / Co ²⁺ / Ca ²⁺	

Question	Answer	Additional Guidance	Mark
number			
1(b)(ii)	An answer that makes reference to the following point:		(1)
	• carbon dioxide / CO ₂ ((g))		

Question	Answer	Additional Guidance	Mark
number			
1(b)(iii)	An answer that makes reference to the following points:		(1)
	• carbonate (ion) / CO ₃ ²⁻ / hydrogencarbonate (ion) / HCO ₃ ⁻	If name and formula are given both must be	
		correct	

Question	Answer		Additional Guidance	Mark
number				
1(b)(iv)	An answer that makes reference to the following points:			(2)
	 (conc) hydrochloric acid / HCl((aq)) (in (b)(i)) the yellow colour is due to CuCl₄²⁻ / tetrachlorocuprate(II) 	(1)(1)	Ignore references to carbonate for HCl Ignore dilute Allow green Allow tetrachlorocopper ion / complex If no other mark is scored allow (1) for any acid justified with the carbonate test	

Question	Answer	Additional Guidance	Mark
number			
1(b)(v)	An answer that makes reference to the following points:		(2)
	• (B) contains a sulfate ion / SO ₄ ²⁻ / CuSO ₄ ((aq)) / copper(II) sulphate (1)	Do not award other compounds containing sulfate ions	
	• (D) is barium chloride (solution) / BaCl ₂ ((aq)) (1)	Allow barium nitrate / Ba(NO ₃) ₂ / Lead(II) nitrate / Pb(NO ₃) ₂	

Question	Answer	Additional Guidance	Mark
number			
1(b)(vi)	An answer that makes reference to the following points:		(2)
	 Copper(II) hydroxide / Cu(OH)₂((s)) / Cu(OH)₂(H₂O)₄((s)) Tetraamminecopper(II) / Tetraamminediaquacopper(II) / [Cu(NH₃)₄(H₂O)₂]²⁺((aq)) / [Cu(NH₃)₄]²⁺((aq)) 	Do not award [Cu(NH ₃) ₆] ²⁺ ((aq)) / hexaaminecopper(II)	

Question	Answer	Additional Guidance	Mark
number 1(b)(vii)	• [CuEDTA] ²⁻ / CuEDTA ²⁻ (1) / Na ₂ [CuEDTA] as a product	Example of equation Allow equations using different starting complex ions or ions such as $[Cu(NH_3)_6]^{2+}/[Cr(NH_3)_6]^{2+}/[Cr(NH_3)_6]^{2+}/[Cr(NH_3)_6]^{2+}/[Cr(NH_3)_6]^{2+}/[Cr(NH_3)_4]^{2+}$ applying TE to the formula and charge of the EDTA complex formed as a product. $Cu^{2+} + EDTA^{4-} \rightarrow CuEDTA^{2-}$ or $Cu^{2+} + Na_4EDTA \rightarrow CuEDTA^{2-} + 4Na^+$ or $[Cu(NH_3)_4(H_2O)_2((aq))]^{2+} + Na_4EDTA \rightarrow [CuEDTA]^{2-} + 4Na^+ + 4NH_3 + 2H_2O$ or $[Cu(NH_3)_4((aq))]^{2+} + EDTA^{4-} \rightarrow [CuEDTA]^{2-} + 4NH_3$	(2)
	• remainder of equation correct (1)	Ignore state symbols even if incorrect	

(Total for Question 1 = 14 marks)

Question number	Answer	Additional Guidance	Mark
2(a)	An answer that makes reference to the following points: • SO ₂ is toxic / corrosive • use a fume cupboard (1)	Assume 'it' is the reaction mixture not specifically SO ₂ Allow just 'SO ₂ (is the main hazard)' Allow SO ₂ is poisonous Ignore pungent/unpleasant smell Dependant on an appropriate hazard being suggested. Allow a well ventilated laboratory Ignore mask Do not award just 'fume cupboard' with no hazard given If M1 is not scored allow M2 for a suitable safety precaution for a stated hazard not associated with SO ₂ e.g. potassium chlorate(V) is corrosive so wear gloves would score M2 If M1 is scored for just 'SO ₂ is the main hazard' do not award M2 If more than one hazard is stated do not allow M1 but allow M2 if one of the hazards has a suitable precaution	(2)

Question	Answer		Additional Guidance	Mark
number				
2(b)			Example of calculation	(2)
	calculation of the number of moles of silver chloride formed	(1)	$= 0.430 \div 143.4 = 0.0029986 / 2.9986 \times 10^{-3} \text{ (mol)}$	
	omondo romida	(1)	= answer to M1 x 10	
	• calculation of the concentration of KClO ₃		$= 0.0029986 \times 10 = 0.029986 / 2.9986 \times 10^{-2} / 0.0300 $ (mol dm ⁻³)	
			Allow TE	
			Do not award incorrect units on the final answer	
			Ignore SF including 1 SF	

Question number	Answer		Additional Guidance	Mark
number 2(c)	An explanation that makes reference to the following points: • silver nitrate/silver nitrate solution is present (because the precipitate is not washed) • (so added mass of silver nitrate/solution) is included in the calculation / mass of AgCl appears higher or	(1)	M2 dependent on the solid containing silver, nitrate solution, water or not being dried properly Allow silver chloride/precipitate is not fully dried Ignore measurement error Allow water in place of silver nitrate solution Allow the mass of AgCl obtained was higher	(2)
	(added mass of silver nitrate / solution) means calculated moles of AgCl / moles of KClO ₃ is higher		Do not award just 'so concentration of KClO ₃ is higher' Do not award just 'mass of AgCl is larger' without a source for the extra mass If M1 is not scored allow one mark for a description of higher calculated mass or moles from other error such as impurities	

Question	Answer	Additional Guidance	Mark
number			
2(d)	An answer that makes reference to the following point:		(1)
	• from green and to yellow	Allow green and to yellow-brown / orange / brown Ignore modifiers e.g. pale Ignore states even if incorrect Do not award red or red in combination with other colours e.g. red-brown	

Question number	Answer		Additional Guidance	Mark
2(e)			Ignore transfer of reaction solution to a conical flask prior to titration Ignore repeating the experiment and averaging results / doing a rough titration then doing an accurate one using the whole reaction mixture	(5)
	• potassium manganate(VII) / KMnO ₄ solution in a burette	(1)	Do not award pipette instead of burette	
	• add (potassium) manganate(VII) drop by drop (at the end-point)	(1)	Ignore transfer slowly	
	• (record the volume when) solution goes (permanent) pink	(1)	Ignore initial colour of titration, even if incorrect e.g. from yellow / colourless / pale green / brown to pink all score (1) Do not award purple	
	additional two suggestions that would improve accuracy of endpoint (1 mark for each)	(2)	e.g. use of a white tile (to more clearly see colour change) rinsing sides of reaction vessel with distilled water swirling/stirring/shaking close to the end-point Rinsing burette with KMnO ₄ solution Read the bottom of the meniscus at eye-level Addition of (dilute) sulfuric acid (added to Fe ²⁺ or KMnO ₄) Do not award concentrated sulfuric acid If a sample is taken from the whole mixture then max 4	
			If titration is reversed (iron(II) in burette) allow TE max 4	

Question	Answer		Additional Guidance	Mark
number 2(f)			Example of calculation:	(6)
	• calculation of initial number of moles of Fe ²⁺ ((1)	$= \frac{150}{1000} \times 0.0750 = 0.01125 / 1.125 \times 10^{-2} $ (mol) (answer 1)	
	• calculation of number of moles of potassium manganate(VII) in titration ((1)	$= \underbrace{9.25}_{1000} \times 0.050 = 0.0004625 / 4.625 \times 10^{-4} \text{ (mol)}$ (answer 2)	
	• calculation of the number of moles of Fe ²⁺ remaining after the reactions	(1)	(answer 2) \times 5 = 0.0023125 / 2.3125 \times 10 ⁻³ (mol) (answer 3)	
	• calculation of the number of moles of Fe ²⁺ that have reacted	(1)	(answer 1) – (answer 3) = $0.01125 - 0.0023125$	
			= $0.01125 - 0.0023125 = 0.0089375 / 8.9375 \times 10^{-3}$ (mol) (answer 4)	
	• calculation of the number of moles of chlorate ions that have reacted ((1)	$\frac{\text{(answer 4)}}{6} = 0.0014895833 / 1.48958 \times 10^{-3} \text{ (mol)}$ 6 (answer 5)	
	• calculation of the concentration of the solution ((1)	= answer $5 \div 0.050 = 0.0297917 / 2.97917 \times 10^{-2}$ (mol dm ⁻³)	
			Allow TE for each stage Ignore SF including 1SF Penalise incorrect units in final mark only Correct answer with no working scores (0)	

Question number	Answer		Additional Guidance	Mark
2(g)	An explanation that makes reference to the following points:		M2 dependent on M1 being scored or an incorrect description of chloride ions / chlorine ions reacting with Fe ²⁺	(2)
	 potassium manganate(VII) reacts with chloride ions to make chlorine / oxidises chloride ions / is a stronger oxidising agent than chlorine or Cl⁻ is a reducing agent (and reduces manganate(VII) ions) 	(1)	Allow just 'chloride ions will react with manganate(VII) ions' Ignore reaction of Fe ²⁺ with chloride ions Do not award 'chlorine ions'	
	• (so the volume of potassium manganate(VII) would increase) and the calculated concentration of potassium chlorate(V) would decrease	(1)	Allow increase in calculated excess Fe ²⁺ , decrease in calculated reacted Fe ²⁺ or calculated moles of ClO ₃ ⁻ . If more than one is given, all must be correct	

(Total for Question 2 = 20 marks)

Question number	Answer	Additional Guidance	Mark
3(a)(i)	An answer that makes reference to the following point:		(1)
	so that the volatile / toxic benzene doesn't escape from the reaction mixture	Must be some mention of benzene Allow the reaction reaches completion / otherwise the reaction would not be complete / to ensure all the benzene reacts Ignore effect on yield	

Question Number	Answer	Additional guidance	Mark
3(a)(ii)		Example of diagram: water out reaction mixture heat source	(3)
	 round-bottomed / pear shaped flask containing mixture and heat (1) vertical condenser with water jacket and water flowing in the correct direction (1) no gaps and open condenser and apparatus would work (1) 	M1 Allow any indication of heat including an arrow or water bath or electrical heater Do not award conical flask/flask with no liquid in Ignore anti bumping granules M3 Ignore thermometer in the top of the condenser if it does not seal the apparatus or through the side of a two necked flask where it is sealed. Ignore stirrer down condenser Do not award if the condenser and flask are one piece of apparatus	
		Allow just M1 for distillation apparatus	

Question	Answer		Additional Guidance	Mark
number				
3(a)(iii)	An answer that makes reference to the following points:			(2)
	reactants are immiscible / form separate layers / do not mix)	Allow are not completely miscible Allow to make the reactants mix	
	(the reactants need to be mixed) to ensure enough contact for a reaction to take place (1))	Allow so allows a reasonable rate of reaction Allow so the reactants come into contact Allow so the reactants are exposed to each other Allow so reaction can take place effectively Allow to ensure even heat distribution Ignore just 'increase the rate of reaction'	

Question	Answer	Additional Guidance	Mark
number			
3(b)	An answer that makes reference to the following points:		(1)
	• (anhydrous) sodium sulfate / Na ₂ SO ₄ / magnesium sulfate / MgSO ₄ / calcium chloride / CaCl ₂	Allow silica gel	
		If both the name and formula are given both must be correct	
		Do not award anhydrous copper sulfate / anhydrous cobalt chloride / conc sulfuric acid	

Question number	Answer	Additional Guidance	Mark
3(c)(i)	An answer that makes reference to the following point:		(1)
	HNO ₂ / nitrous acid is unstable / decomposes (so cannot be transported)	Ignore difficult to store Do not award reacts to form nitric acid	

Question number	Answer		Additional Guidance	Mark
3(c)(ii)	An explanation that makes reference to the following points:			(2)
	• below 0°C the reaction is too slow	(1)	Allow reaction mixture may solidify Allow less energy for overcoming the activation barrier Ignore reaction will stop / does not occur Do not award not enough energy to overcome the activation barrier	
	above 10°C the diazonium chloride / compound / ion decomposes / hydrolyses / reacts with water / forms phenol (and nitrogen gas)	(1)	Allow phenol is formed Allow the product decomposes Allow phenylamine reacts to form phenol Ignore nitrous acid decomposes Ignore it decomposes Ignore waste products are formed Do not award the azo dye / Organol Brown / phenylamine would decompose Do not award multiple substitutions will occur	

Question	Answer	Additional Guidance	Mark
number			
3(d)(i)	An answer that makes reference to the following point:		(1)
	 (so a lot of the azo dye) does not remain in solution (when it cools) or gives a saturated solution (when it has cooled) 	Ignore 'to obtain a concentrated solution' Do not award just to maximize yield	

Question	Answer	Additional Guidance	Mark
number			
3(d)(ii)	An answer that makes reference to the following point:		(1)
	• (pre-heated) to prevent (premature) crystallisation (of the azo dye in the funnel)	Allow to keep the Organol Brown / product / solid in solution Ignore it remains in the liquid state	

Question	Answer	Additional Guidance	Mark
number			
3(d)(iii)	An answer that makes reference to the following points:		(2)
	• Step 2 is to remove insoluble impurities (1)	If no other mark is scored just 'to remove insoluble	
		and soluble impurities' scores (1)	
	• Step 3 is to remove soluble impurities (1)		
		Do not award 'to remove soluble and insoluble	
		impurities'	

Question	Answer	Additional Guidance	Mark
number			
3(d)(iv)	An answer that makes reference to the following point:		(1)
	 to avoid (thermal) decomposition 	Allow to avoid melting the dye / crystals	
		Ignore relative speed of drying	
		Do not award to stop it making a solution	
		Do not award decay	

Question	Answer	Additional Guidance	Mark
number			
3(e)	An answer that makes reference to the following point:		(1)
	sharp melting temperature	Allow any indication of a small range (±2°C) Ignore melting temperature matching data in a data book Ignore impurities make the melting point lower / higher than the literature value	

(Total for Question 3 = 16 marks) Total for Paper = 50 marks