



Mark Scheme (Results)

January 2020

Pearson Edexcel International Advanced
Subsidiary Level In Physics (WPH11)


Paper 01 Mechanics and Materials

Question Number	Answer	Mark
1	<p>C is the correct answer</p> <p>A is not the correct answer as units would be kg m s^{-2} B is not the correct answer as units would be kg m s^{-1} D is not the correct answer as units would be $\text{kg m}^2 \text{s}^{-2}$</p>	(1)
2	<p>B is the correct answer</p> <p>A is not the correct answer as force per unit length has no meaning. C is not the correct answer as this is the gravitational force. D is not the correct answer as this is gravitational potential.</p>	(1)
3	<p>B is the correct answer</p> <p>A is not the correct answer as the magnitude of the final velocity would be greater and a little less to the right. C is not the correct answer as the final velocity would be greater and to the left of the original velocity. D is not the correct answer as the final velocity would be similar to C, but more to the left and of a lesser magnitude.</p>	(1)
4	<p>D is the correct answer</p> <p>A is not the correct answer as the velocity is not constant at all times. B is not the correct answer as the velocity is still not constant at all times. C is not the correct answer as the air resistance does not act in the opposite direction to gravity when an object travels upwards.</p>	(1)
5	<p>D is the correct answer</p> <p>A is not the correct answer as the areas under the two graphs are not both zero. B is not the correct answer as the areas under the two graphs are not equal. C is not the correct answer as P is <u>at</u> the initial position.</p>	(1)
6	<p>A is the correct answer</p> <p>B is not the correct answer as the change in velocity is not 2 m s^{-1}. C is not the correct answer as it gives a negative time, and the collision takes a positive amount of time. D is not the correct answer for the same reason that B is not.</p>	(1)
7	<p>D is the correct answer</p> <p>A is not the correct answer as it contradicts Newton's Third Law. B is not the correct answer as it also contradicts Newton's Third Law. C is not the correct answer as the force of Y on X is in the opposite direction to the velocity of X.</p>	(1)
8	<p>D is the correct answer</p> <p>A is not the correct answer as it gives units of J^{-1} which is not a unit for energy. B is not the correct answer for the same reason that A is not. C is not the correct answer as $68 \neq 68\%$.</p>	(1)
9	<p>C is the correct answer</p> <p>A is not the correct answer as every column is wrong. B is not the correct answer as the P and Q columns are the wrong way round. D is not the correct answer as the Q and R columns are the wrong way round.</p>	(1)

10	<p>A is the correct answer</p> <p>B is not the correct answer as the gravitational force does not increase. C is not the correct answer as an increase in temperature would reduce the viscosity which would not account for a decrease in acceleration. D is not the correct answer as the upthrust depends on the density of the fluid and the volume of the sphere, neither of which changes.</p>	(1)
-----------	---	------------

Question Number	Answer	Mark
11	<ul style="list-style-type: none"> • Reference to $s = ut + \frac{1}{2}at^2$ with $u = 0$ (1) • Correct variable labels on graph axes to give a straight line through origin. (1) • Reference to time in s and distance in m (this can be taken from the axes labels or a suitable unit conversion) (1) • Straight line through origin. (1) • Correct method to determine g using their graph. (1) 	(5)
	Total for question 11	5

Question Number	Answer	Mark
12(a)	<ul style="list-style-type: none"> Laminar/non-turbulent flow Or Slow moving sphere 	(1) (1)
12(b)	<ul style="list-style-type: none"> Use of $W = mg$ $W = U + D$ Use of $F = 6\pi r\eta v$ Use of $W = U + D$ to obtain quantity to compare, e.g. $D = (-) 7.8 \times 10^{-3} \text{ (N)}$ Comparison leading to valid conclusion from candidate's calculation. e.g. $F = 2.5 \times 10^{-5} \text{ N} \neq D$ or $v = 155 \text{ m s}^{-1} \neq 0.5 \text{ m s}^{-1}$ <i>et al.</i> <p><u>Example of calculation</u> $W = 9.1 \times 10^{-4} \text{ kg} \times 9.81 \text{ N kg}^{-1} = 9.0 \times 10^{-3} \text{ N}$ $\Sigma F = 9.0 \times 10^{-3} \text{ N} - 1.1 \times 10^{-3} \text{ N} - \text{drag} = 0$ Drag = (-) $7.9 \times 10^{-3} \text{ N}$ $F = 6 \times \pi \times 3.0 \times 10^{-3} \text{ m} \times 8.9 \times 10^{-4} \text{ Pa s} \times 0.50 \text{ m s}^{-1}$ $F = 2.5 \times 10^{-5} \text{ N}$</p>	(1) (1) (1) (1) (1) (5)
	Total for question 12	6

Question Number	Answer	Mark
13(a)	<ul style="list-style-type: none"> Point at which weight is taken to act. 	(1)
		(1)
13(b)	<ul style="list-style-type: none"> Gradient everywhere positive Stops at $R = \text{weight}/W/mg$ at $l/2$ Starts at $R = \frac{1}{2} \text{ weight}/0.5W/0.5mg$ etc. 	(1) (1) (1)
		(3)
13(c)	<ul style="list-style-type: none"> Centre of gravity/mass is not above the shelf. Or Line of action of weight does not pass through the shelf. There is a net moment clockwise. Or No anticlockwise moment to balance moment of weight. 	(1) (1)
		(2)
	Total for question 13	6

Question Number	Answer	Mark																																								
*14	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content and lines of reasoning.</p> <table><tr><th>IC points</th><th>IC mark</th><th>Max linkage mark available</th><th>Max final mark</th></tr><tr><td>6</td><td>4</td><td>2</td><td>6</td></tr><tr><td>5</td><td>3</td><td>2</td><td>5</td></tr><tr><td>4</td><td>3</td><td>1</td><td>4</td></tr><tr><td>3</td><td>2</td><td>1</td><td>3</td></tr><tr><td>2</td><td>2</td><td>0</td><td>2</td></tr><tr><td>1</td><td>1</td><td>0</td><td>1</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> <table><tr><td></td><td>Number of marks awarded for structure of answer and sustained line of reasoning</td></tr><tr><td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout</td><td>2</td></tr><tr><td>Answer is partially structured with some linkages and lines of reasoning</td><td>1</td></tr><tr><td>Answer has no linkages between points and is unstructured</td><td>0</td></tr></table> <p>Indicative content</p> <ul style="list-style-type: none">No change in properties at low strain/stress. Or A change in properties after a certain point/stress.Energy absorbed by capsule concrete can be greater than plain concrete.Area under graph for capsule concrete greater.(At greater stress) capsule concrete is stiffer Or (at greater stress) capsule concrete has greater Young Modulus.Graph is steeper for capsule concrete (at high stress) Or ratio of stress to strain is greater (at high stress).Maximum stress/force greater for capsule concrete Or capsule concrete is stronger.	IC points	IC mark	Max linkage mark available	Max final mark	6	4	2	6	5	3	2	5	4	3	1	4	3	2	1	3	2	2	0	2	1	1	0	1	0	0	0	0		Number of marks awarded for structure of answer and sustained line of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2	Answer is partially structured with some linkages and lines of reasoning	1	Answer has no linkages between points and is unstructured	0	(6)
IC points	IC mark	Max linkage mark available	Max final mark																																							
6	4	2	6																																							
5	3	2	5																																							
4	3	1	4																																							
3	2	1	3																																							
2	2	0	2																																							
1	1	0	1																																							
0	0	0	0																																							
	Number of marks awarded for structure of answer and sustained line of reasoning																																									
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2																																									
Answer is partially structured with some linkages and lines of reasoning	1																																									
Answer has no linkages between points and is unstructured	0																																									
	Total for question 14	6																																								

Question Number	Answer	Mark
15(a)	<ul style="list-style-type: none"> Use of $v^2 = u^2 + 2as$ AND $u = 0$ (1) Or $mgh = \frac{1}{2}mv^2$ $v = 3.4 \text{ (m s}^{-1}\text{)}$ (1) <p><u>Example of calculation</u> $v^2 = 2 \times 9.81 \text{ m s}^{-2} \times 0.60 \text{ m}$ $v = \sqrt{11.77} \text{ m s}^{-1}$ $v = 3.43 \text{ m s}^{-1}$</p>	(2)
15(b)	<ul style="list-style-type: none"> Horizontal $3.4 \times \sin 70^\circ$ Or $3.4 \times \cos 20^\circ$ Or calculated value. (1) Vertical $3.4 \times \cos 70^\circ$ Or $3.4 \times \sin 20^\circ$ 1.16 Or calculated value. (1) <p>Allow e.c.f. from part (a)</p>	(2)

15(c)	<p>Use of $v = s/t$ to determine time to end of ramp (0.38 s).</p> <ul style="list-style-type: none"> Use of $s = ut - \frac{1}{2} g t^2$ to determine drop in altitude after time t (0.27 m). (1) Ball does not bounce on the ramp. (1) Justifies conclusion from numbers calculated. e.g. $0.86 - 0.27 > 0.00$ means has not reached ground by end of ramp. (1) <p><u>Example of calculation</u> (1)</p> $t = \frac{1.23 \text{ m}}{3.4 \text{ m s}^{-1} \times \sin 70^\circ}$ $t = 0.39 \text{ s}$ $s = (3.4 \text{ (m s}^{-1}) \times \cos 70^\circ \times 0.39 \text{ s}) + (\frac{1}{2} \times (-9.81 \text{ m s}^{-2}) \times (0.39 \text{ s})^2)$ $s = -0.28 \text{ m}$ <p>Or</p> <ul style="list-style-type: none"> Use of $s = ut - \frac{1}{2} g t^2$ to determine time to $s = -0.86$ Use of $s = vt$ to calculate s (1) Ball does not bounce on the ramp. (1) Justifies conclusion from numbers calculated. e.g. $1.23 < 1.79$ (1) <p>Or (1)</p> <ul style="list-style-type: none"> Use of $v = u - gt_1$ with $v = 0$ to get time to max height (0.12s) and use of $s = \frac{1}{2} g t_1^2$ to get gain in height (0.07m) and use of $s = ut + \frac{1}{2} g t_2^2$ with $u = 0$ and $s = 0.93$ to get time from there to the ground (0.44s) Total time $t = t_1 + t_2$. (1) Use of $s = vt$ to calculate s (1) Ball does not bounce on the ramp. (1) Justifies conclusion from numbers calculated. e.g. $1.23 < 1.79$ (1) 	(4)
	Total for question 15	8

Question Number	Answer	Mark
16(a)(i)	<ul style="list-style-type: none"> Use measurement and scaling factor (1) (1) $h = 3.4 \pm 0.1$ cm (1) <p>OR</p> <ul style="list-style-type: none"> Use of $s = \frac{1}{2}at^2$ with $t = 0.083$ s (1) $h = 3.4$ cm (1) 	(2)
16(a)(ii)	<ul style="list-style-type: none"> Use of (average) speed = s/t with s = value from part (a)(i) (1) Use of initial speed = $2 \times$ average speed (1) Speed = 0.82 m s^{-1} ecf from (a)(i) (1) <p>OR</p> <ul style="list-style-type: none"> Use of $E_g = mgh$ with h = value from (a)(i) (1) Use of $\frac{1}{2}mv^2 = E_g(\text{initial})$ (1) Speed = 0.82 m s^{-1} ecf from (a)(i) (1) <p>OR</p> <ul style="list-style-type: none"> Use of $v = u + at$ with $v = 0$ (1) Use of $a = -g$ (1) Speed = $9.81 \times 0.083 = 0.81 \text{ m s}^{-1}$ (1) <p>OR</p> <ul style="list-style-type: none"> Use of with $s = ut + \frac{1}{2}at^2$ with s = value from part (a)(i) (1) Use of $a = -g$ (1) Speed = 0.82 m s^{-1} ecf from (a)(i) (1) <p><u>Example of calculation</u> $u = s/t - \frac{1}{2}at$ $u = 0.034/0.083 + \frac{1}{2} \times 9.81 \times 0.083$</p> <p>OR</p> <ul style="list-style-type: none"> Use of $v^2 = u^2 + 2as$ with s = value from (a)(i) and $v = 0$ (1) Use of $a = -g$ (1) 	

	<ul style="list-style-type: none"> Speed = 0.82 m s^{-1} ecf from (a)(i) (1) <p><u>Example of calculation</u></p> <p>Actual distance travelled by popcorn = $6.2 \text{ cm} \div 1.8 = 3.4 \text{ cm}$</p> <p>(average speed) = $\frac{0.034 \text{ m}}{83 \times 10^{-3} \text{ s}} = 0.41 \text{ m s}^{-1}$</p> <p>Initial speed = $2 \times \text{average speed} = 0.82 \text{ m s}^{-1}$ (3)</p>	
16(b)	<ul style="list-style-type: none"> Use of 14% to determine mass (of water/popcorn). (1) <p>Or</p> <p>$m_{\text{popcorn}}/m_{\text{water}} = 86/14$</p> <ul style="list-style-type: none"> Use of $p = mv$ (1) Use of momentum conservation (1) $v = (-) 9.2 \text{ m s}^{-1}$ (1) <p><u>Example of calculation</u></p> <p>$0 = (0.0946 \text{ g} \times 1.5 \text{ m s}^{-1}) + (0.0154 \text{ g} \times v)$</p> <p>$v = \frac{-0.0946 \text{ g} \times 1.5 \text{ m s}^{-1}}{0.0154 \text{ g}}$</p> <p>$v = -9.21 \text{ m s}^{-1}$</p> <p>(4)</p>	
Total for question 16		9

Question Number	Answer	Mark
17(a)(i)	<ul style="list-style-type: none"> • $E_{el} = \frac{1}{2} k \Delta x^2$ Or Use of $E_{el} = \frac{1}{2} F \Delta x$ and use of $F = k \Delta x$. (1) • Elastic PE is transferred into kinetic energy Or $E_{el} = E_k$ (1) • $\frac{1}{2} m v^2 = \frac{1}{2} k \Delta x^2$ (1) • States that m and k are constant so $v \propto \Delta x$. Or States that $= \sqrt{\frac{k}{m}} \Delta x$. (1) 	(4)
17(a)(ii)	<ul style="list-style-type: none"> • Gradient calculated. Or Use of a point on the line in a relevant equation. (1) • Use of $\frac{1}{2} k \Delta x^2 = \frac{1}{2} m v^2$ or gradient $= \sqrt{(k/m)}$ i.e. $k = m \times \text{gradient}^2$ (1) • k in range 22 – 26 N m⁻¹ (1) <p><u>Example of calculation</u> Gradient $\frac{4.8 \text{ m s}^{-1} - 2.2 \text{ m s}^{-1}}{0.30 \text{ m}} = 8.67 \text{ (s}^{-1}\text{)}$ $k = \text{mass} \times \text{gradient}^2$ $k = 3.0 \times 10^{-1} \text{ kg} \times (8.67 \text{ s}^{-1})^2$ $k = 22.6 \text{ N m}^{-1}$</p>	(3)
17(b)	<ul style="list-style-type: none"> • Limit of proportionality exceeded. Or Extension no longer proportional to force. (1) • Range of Hooke's Law exceeded. Or Hooke's Law no longer applies. (1) 	(2)
	Total for question 17	9

Question Number	Answer	Mark
18(a)	<ul style="list-style-type: none"> Ratio of stress to strain (for a material). Or stress per unit strain. Or σ / ϵ with symbols defined. Or $\frac{F x}{A \Delta x}$ with symbols defined. 	(1) (1)
18(b)(i)	<ul style="list-style-type: none"> Mean diameter = 0.234 mm (rounds to) Use of $A = \pi r^2$ $A = 4.3 \times 10^{-8} \text{ m}^2$ or 0.043 mm² <p><u>Example of calculation</u> Mean diameter = $\frac{1}{4} (0.230 + 0.235 + 0.230 + 0.240) = 0.234 \text{ mm}$ Area = $\pi \frac{(0.234 \times 10^{-3} \text{ m})^2}{4} = 4.30 \times 10^{-8} \text{ m}^2$</p>	(1) (1) (1) (3)
18(b)(ii)	<ul style="list-style-type: none"> Use of $W = m g$ Use of gradient = $m / \Delta x$ in Young Modulus formula i.e. $E = \text{gradient} \times g \times x / A$ $E = 1.6 \times 10^{11} \text{ Pa}$ e.c.f. from (b)(i) <p><u>Example of calculation</u> Young modulus = $195 \times 9.81 \text{ N kg}^{-1} \times \frac{3.50 \text{ m}}{4.30 \times 10^{-8} \text{ m}^2}$ $= 1.56 \times 10^{11} \text{ Pa}$</p>	(1) (1) (1) (3)
18(b)(iii)	<p>Shorter wire gives greater gradient.</p> <p>Young modulus the same.</p>	(1) (1) (2)
	Total for question 18	9

Question Number	Answer	Mark
19(a)	<ul style="list-style-type: none"> Upthrust/U upwards (1) Tension/T upwards (1) Weight/W/mg downwards (1) <p>(-1 for each extra force over three, -1 if any arrow does not touch the dot, -1 if any arrow is not close to vertical. Accept a single line up with two labelled arrowheads.)</p>	(3)
19(b)(i)	<ul style="list-style-type: none"> Water exerts upward force on sphere. (1) Or Water exerts an upthrust on the sphere. Sphere exerts a downwards/opposite force on water by Newton's Third Law. (1) Extra downward force on water (increases reading on balance). (1) 	(3)
19(b)(ii)	<ul style="list-style-type: none"> Mass of displaced water = 150 g (1) Use of $V = m/\rho$ for water with $\rho = 1\,000\text{ kg m}^{-3}$ (150 ml) (1) Use of $m = \rho V$ for sphere WITH $\rho = 2\,000\text{ kg m}^{-3}$ (1) $m = 0.30\text{ kg}$ (1) <p><u>Example of calculation</u> Increase in weight of water = force of ball on water = upthrust on ball Mass of displaced water = $465\text{ g} - 315\text{ g} = 150\text{ g}$ Volume of sphere = $\frac{0.150\text{ kg} \times g}{1000\text{ kg m}^{-3} \times g} = 1.5 \times 10^{-4}\text{ m}^3$ Mass of sphere = $2000\text{ kg m}^{-3} \times 1.5 \times 10^{-4}$</p>	(4)
19(b)(iii)	<ul style="list-style-type: none"> Upthrust less in oil or weight of displaced oil less or downward force of sphere on oil less (1) (Therefore increase in) balance reading less (than for water). (1) <p>(MP2 dependent on MP1)</p>	(2)
Total for question 19		12