



Mark Scheme (Results)

January 2022

Pearson Edexcel International Advanced
Subsidiary Level In Physics (WPH15) Paper 01
Thermodynamics, Radiation, Oscillations and
Cosmology

Question Number	Answer	Mark
1	B is the correct answer , as this is part of the definition of s.h.m.	(1)
2	B is the correct answer A is incorrect, as this would increase the value of L_V C is incorrect, as it's not necessary to stir boiling water D is incorrect, as this would increase the value of L_V	(1)
3	D is the correct answer , as $T = 2\pi\sqrt{\frac{L}{g}}$ and $T = 2 \times 8.25\text{s}$	(1)
4	C is the correct answer , as the activity halves in each half-life period	(1)
5	A is the correct answer , as $pV \propto T$	(1)
6	D is the correct answer , as this is a statement of Hubble's law	(1)
7	B is the correct answer A is incorrect, as background count rate varies from place to place C is incorrect, as the background count is not constant D is incorrect, as some detector are more sensitive than others	(1)
8	B is the correct answer , as $I = \frac{L}{4\pi d^2}$	(1)
9	B is the correct answer A is incorrect, as the lines may be shifted into any region of the spectrum C is incorrect, as the intensity of the lines is not related to the redshift D is incorrect, as the wavelengths of the emitted lines is not affected	(1)
10	D is the correct answer , A is incorrect, as ^{56}Fe is the most stable isotope B is incorrect, as the graph shows the binding energy per nucleon C is incorrect, as high mass nuclei could be fused as long as energy is supplied	(1)

Question Number	Answer	Mark
11	At least 1 cycle of a sinusoidal graph Displacement axis shows amplitude as 5 cm Use of $a = (-)\omega^2 x$ and $\omega = \frac{2\pi}{T}$ to calculate T Time axis shows period as calculated value of T <u>Example of calculation</u> $\omega = \sqrt{\frac{8.0 \text{ cm s}^{-2}}{5.0 \text{ cm}}} = 1.26 \text{ s}^{-1}$ $T = \frac{2\pi}{1.26 \text{ s}^{-1}} = 4.97 \text{ s}$	(1) (1) (1) (1) 4
Total for question 11		4

Question Number	Answer	Mark
12	<p>Energy transferred from hot liquid = energy transferred to ice + energy transferred to cold water (1)</p> <p>[This may be implicit]</p> <p>Use of $E = mc\Delta\theta$ (1)</p> <p>Use of $E = mL$ (1)</p> <p>Mass of ice required to cool drink to 58 °C is 2.4×10^{-2} kg Or Final temperature using 4 g of ice is 69 °C (1)</p> <p>Valid conclusion based on a consideration of their calculated value in comparison with a corresponding value in the question. (1)</p> <p><u>Example of calculation</u></p> <p>Energy transferred from hot liquid = energy transferred to ice + energy transferred to cold water</p> <p>$0.275 \text{ kg} \times 3750 \text{ J kg}^{-1}\text{K}^{-1} \times (71.5 - 58.0)\text{K}$ $= m \times 3.34 \times 10^5 \text{ J kg}^{-1} + m \times 4190 \text{ J kg}^{-1}\text{K}^{-1} \times (58.0 - 0)\text{K}$ $\therefore 1.39 \times 10^4 \text{ J} = m \times (3.34 \times 10^5 + 2.43 \times 10^5) \text{ J kg}^{-1}$ $\therefore m = \frac{1.39 \times 10^4 \text{ J}}{5.77 \times 10^5 \text{ J kg}^{-1}} = 2.41 \times 10^{-2} \text{ kg} = 24 \text{ g}$ So 4 g would not bring the temperature below the ideal serving temperature.</p>	5
	Total for question 12	5

Question Number	Answer	Mark
13(a)	<p>Use of $pV = NkT$ (1)</p> <p>Temperature converted to kelvin (1)</p> <p>$V = 6.9 \text{ m}^3$ (1)</p> <p><u>Example of calculation</u></p> <p>$\frac{pV}{T} = \text{a constant}$</p> $\frac{8.4 \times 10^4 \text{ Pa} \times V}{(273 - 48) \text{ K}} = \frac{1.02 \times 10^5 \text{ Pa} \times 7.50 \text{ m}^3}{(273 + 22.5) \text{ K}}$ $\therefore V = \frac{1.02 \times 10^5 \text{ Pa} \times 7.5 \text{ m}^3 \times (273 - 48) \text{ K}}{(273 + 22.5) \text{ K} \times 8.4 \times 10^4 \text{ Pa}} = 6.93 \text{ m}^3$	3
13(b)	<p>Use of $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$ (1)</p> <p>Decrease = $1.5 \times 10^{-21} \text{ J}$ (1)</p> <p><u>Example of calculation</u></p> $\Delta(\text{mean kinetic energy}) = \frac{3}{2} 1.38 \times 10^{-2} \text{ J K}^{-1}(-48 - 22.5) \text{ K}$ $\therefore \Delta(\text{mean kinetic energy}) = -1.46 \times 10^{-21} \text{ J}$	2
Total for question 13		5

Question Number	Answer	Mark
14	Max kinetic energy read from graph	(1)
	Use of 15.6 eV to calculate number of nitrogen molecules ionised	(1)
	Use of 250 to calculate range of β particle	(1)
	Range of β particle read from graph	(1)
	Comparison of their two ranges with conclusion	(1)
	OR	
	Max kinetic energy read from graph	(1)
	Use of 15.6 eV to calculate number of nitrogen molecules ionised	(1)
	Range of β particle read from graph	(1)
	Use of range to calculate number of molecules ionised	(1)
	Comparison of their two numbers of molecules with conclusion	(1)
	<u>Example of calculation</u>	
	Maximum $E_k = 0.52 \text{ MeV} \rightarrow 0.55 \text{ MeV}$	
	$N = \frac{5.3 \times 10^5 \text{ eV}}{15.6 \text{ eV}} = 3.40 \times 10^4$	
	$\text{Range} = \frac{3.40 \times 10^4}{250 \text{ cm}^{-1}} = 136 \text{ cm} = 1.36 \text{ m}$	
	Range of β particle = 1.2 m \rightarrow 1.4 m	
Total for question 14		5



Question Number	Answer	Mark
15	<p>λ_{\max} read from graph [450 nm \rightarrow 500 nm] (1)</p> <p>Use of $\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m K}$ (1)</p> <p>Use of $L = \sigma AT^4$ (1)</p> <p>Use of $A = 4\pi r^2$ (1)</p> <p>$D_P = 4.6 \times D_{\text{Sun}}$ so statement is incorrect (1)</p> <p>Or $D_P = 3.2 \times 10^9 \text{ m}$, which is more than twice Sun's diameter, so statement is incorrect</p> <p><u>Example of calculation</u></p> $T = \frac{2.898 \times 10^{-3} \text{ m K}}{470 \times 10^{-9} \text{ nm}} = 6170 \text{ K}$ $A = \frac{2.65 \times 10^{27} \text{ W}}{5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^4 \times (6170 \text{ K})^4} = 3.22 \times 10^{19} \text{ m}^2$ $r = \sqrt{\frac{3.22 \times 10^{19} \text{ m}^2}{4\pi}} = 1.60 \times 10^9 \text{ m}$ $\frac{D_P}{D_{\text{Sun}}} = \frac{2 \times 1.60 \times 10^9 \text{ m}}{6.96 \times 10^8 \text{ m}} = 4.6$	5
	Total for question 15	5

Question Number	Answer	Mark
16(a)(i)	$v \propto \sqrt{\frac{M}{r}}$ <p>(1)</p> <p>Within the central region M changes a lot (so v increases) Or Outside the central region M is approximately constant (so v decreases)</p> <p>As r increases v reaches a peak value as shown on the graph (1)</p> <p>[A bald description of the graph having a peak value can score MP3] (1)</p>	3
16(a)(ii)	<p>There must be more mass (than we can observe) (1)</p> <p>[Accept statement that there must be a greater gravitational force] (1)</p> <p>There is dark matter present (in the galaxy)</p>	2
16(b)	<p>(For a closed universe) the density of the universe must be greater than the critical density (1)</p> <p>And the (average) density of the universe is uncertain (1)</p> <p>Or the amount of dark matter is uncertain</p>	2
	Total for question 16	7

Question Number	Answer	Mark
17(a)	<p>Either</p> <p>Current carrying coil/conductor in a magnetic field (1)</p> <p>Coil experiences a force (1)</p> <p>Force changes direction with current (as current is changing direction) (1)</p> <p>Or</p> <p>Current in coil causes a magnetic field (1)</p> <p>Field interacts with permanent magnet's field, so force on coil (1)</p> <p>Field changes direction with current so force changes direction (1)</p>	3
17(b)(i)	<p>Use of $\omega = 2\pi f$ (1)</p> <p>Use of $v = -A\omega \sin \omega t$ (1)</p> <p>$v = 0.82 \text{ m s}^{-1}$ (1)</p> <p><u>Example of calculation</u></p> <p>$\omega = 2\pi \text{ rad} \times 75 \text{ s}^{-1} = 471 \text{ rad s}^{-1}$</p> <p>$v = 1.75 \times 10^{-3} \text{ m} \times 471 \text{ s}^{-1} \times 1 = 0.8247 \text{ m s}^{-1}$</p>	3
17(b)(ii)	At the equilibrium/undisplaced/central/middle (position) (1)	1
17(c)	<p>MAX 2</p> <p>The driver frequency of the coil matches the natural frequency of the cone (1)</p> <p>There is a maximum transfer of energy (from the coil to the cone) (1)</p> <p>Resonance occurs (1)</p> <p>[For full marks the response must be related to the question context]</p>	2
	Total for question 17	9

Question Number	Answer	Mark
18(a)	<p>Use of $T^2 = KR^3$ (1)</p> <p>K for Earth = $2.96 \times 10^{-19} \text{ (s}^2 \text{ m}^{-3}\text{)}$ (1)</p> <p>K for Mars = $2.97 \times 10^{-19} \text{ (s}^2 \text{ m}^{-3}\text{)}$ (1)</p> <p><u>Example of calculation</u></p> $K = \frac{T^2}{R^3} = \frac{(3.16 \times 10^7 \text{ s})^2}{(1.50 \times 10^{11} \text{ m})^3} = 2.959 \times 10^{-19} \text{ s}^2 \text{ m}^{-3}$ $K = \frac{T^2}{R^3} = \frac{(5.93 \times 10^7 \text{ s})^2}{(2.28 \times 10^{11} \text{ m})^3} = 2.967 \times 10^{-19} \text{ s}^2 \text{ m}^{-3}$	3
18(b)	<p>Either</p> <p>Use of $F = \frac{GMm}{r^2}$ with $F = \frac{mv^2}{r}$ (1)</p> <p>Re-arrangement with $v = \frac{2\pi r}{T}$ to identify K as $\frac{(2\pi)^2}{GM}$ (1)</p> <p>$K = 2.97 \times 10^{-19} \text{ (s}^2 \text{ m}^{-3}\text{)}$ (1)</p> <p>Or</p> <p>Use of $F = \frac{GMm}{r^2}$ with $F = m\omega^2 r$ (1)</p> <p>Re-arrangement with $\omega = \frac{2\pi}{T}$ to identify K as $\frac{(2\pi)^2}{GM}$ (1)</p> <p>$K = 2.97 \times 10^{-19} \text{ (s}^2 \text{ m}^{-3}\text{)}$ (1)</p> <p><u>Example of calculation</u></p> $\frac{GMm}{r^2} = m\omega^2 r$ $\frac{GM}{r^2} = \left(\frac{2\pi}{T}\right)^2 r$ $T^2 = \frac{(2\pi)^2}{GM} r^3$ $K = \left(\frac{4\pi^2}{6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2} \times 1.99 \times 10^{30} \text{ kg}}\right) = 2.97 \times 10^{-19} \text{ s}^2 \text{ m}^{-3}$	3

18(c)	Use of $T^2 = KR^3$ (1)	
	$T = 43$ hours (1)	2
	<u>Example of calculation</u> $\left(\frac{T_I}{T_G}\right)^2 = \left(\frac{R_I}{R_G}\right)^3$ $T = \sqrt{\left(\frac{4.22 \times 10^8 \text{ m}}{1.07 \times 10^9 \text{ m}}\right)^3 \times (172 \text{ hour})^2} = 42.6 \text{ hours}$	
	Total for question 18	8

Question Number	Answer	Mark
19(a)	<p>Use of trigonometry to calculate the parallax angle Or Use of trigonometry to calculate distance (1)</p> <p>(Smallest) parallax angle = 3.3×10^{-7} (rad) Or max distance = 6.25×10^{17} (m) (1)</p> <p>Comparison of calculated value with corresponding value in question with valid conclusion (1)</p> <p><u>Example of calculation</u></p> $\sin \alpha = \frac{1.5 \times 10^{11} \text{ m}}{d}$ $\alpha = \sin^{-1} \left(\frac{1.5 \times 10^{11} \text{ m}}{4.6 \times 10^{17} \text{ m}} \right) = 3.26 \times 10^{-7} \text{ rad}$ <p>Or $\alpha = \left(\frac{1.5 \times 10^{11} \text{ m}}{4.6 \times 10^{17} \text{ m}} \right) = 3.26 \times 10^{-7} \text{ rad}$ (small angle approximation)</p> 	3
19(b)	<p>The intensity (of radiation from the candle) is measured (1)</p> <p>The luminosity of the standard candle is known (1)</p> <p>The inverse square law is used to determine the distance [Accept reference to $I=L/4\pi d^2$ with symbols defined] (1)</p>	3
19(c)(i)	<p>Axis labelled with T / K (1)</p> <p>Reverse logarithmic scale (1)</p> <p>6000 K in correct position on scale (1)</p> <p><u>Example of graph labelling</u></p> 	3

19(c)(ii)	<table><tr><th>Description</th><th>Zone</th></tr><tr><td>High mass hot stars</td><td>Z1</td></tr><tr><td>Low mass cool stars</td><td>Z5</td></tr><tr><td>Low mass hot stars</td><td>Z2</td></tr></table>	Description	Zone	High mass hot stars	Z1	Low mass cool stars	Z5	Low mass hot stars	Z2	(1) (1) (1)	3																																
Description	Zone																																										
High mass hot stars	Z1																																										
Low mass cool stars	Z5																																										
Low mass hot stars	Z2																																										
19(c)(iii)	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table><tr><td></td><td>Number of marks awarded for structure of answer and sustained line of reasoning</td></tr><tr><td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout</td><td>2</td></tr><tr><td>Answer is partially structured with some linkages and lines of reasoning</td><td>1</td></tr><tr><td>Answer has no linkages between points and is unstructured</td><td>0</td></tr></table> <p>Total marks awarded is the sum of marks for indicative content and the marks for structure and lines of reasoning</p> <table><tr><td>IC points</td><td>IC mark</td><td>Max linkage mark</td><td>Max final mark</td></tr><tr><td>6</td><td>4</td><td>2</td><td>6</td></tr><tr><td>5</td><td>3</td><td>2</td><td>5</td></tr><tr><td>4</td><td>3</td><td>1</td><td>4</td></tr><tr><td>3</td><td>2</td><td>1</td><td>3</td></tr><tr><td>2</td><td>2</td><td>0</td><td>2</td></tr><tr><td>1</td><td>1</td><td>0</td><td>1</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> <p>Indicative content</p> <p>IC1 The star is fusing hydrogen in its core</p> <p>IC2 When fusion ceases (the core of the star cools and) the core collapses/contracts (under gravitational forces)</p> <p>IC3 The star (moves to Z4 as it expands and) becomes a red giant star</p> <p>IC4 Temperature (in the core) is high enough for helium fusion to begin</p> <p>IC5 Helium begins to run out and then fusion ceases</p> <p>IC6 The star becomes a white dwarf (in Z2)</p>		Number of marks awarded for structure of answer and sustained line of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2	Answer is partially structured with some linkages and lines of reasoning	1	Answer has no linkages between points and is unstructured	0	IC points	IC mark	Max linkage mark	Max final mark	6	4	2	6	5	3	2	5	4	3	1	4	3	2	1	3	2	2	0	2	1	1	0	1	0	0	0	0		6
	Number of marks awarded for structure of answer and sustained line of reasoning																																										
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2																																										
Answer is partially structured with some linkages and lines of reasoning	1																																										
Answer has no linkages between points and is unstructured	0																																										
IC points	IC mark	Max linkage mark	Max final mark																																								
6	4	2	6																																								
5	3	2	5																																								
4	3	1	4																																								
3	2	1	3																																								
2	2	0	2																																								
1	1	0	1																																								
0	0	0	0																																								
Total for question 18			18																																								

Question Number	Answer	Mark
20(a)	<p>Top row correct (1)</p> <p>Bottom row correct (1)</p> <p><u>Example of calculation</u></p> ${}^{225}_{89}\text{Ac} \rightarrow {}^{221}_{87}\text{Fr} + {}^4_2\alpha$	2
20(b)	<p>Use of $1 \text{ u} = 1.66 \times 10^{-27} \text{ kg}$ (1)</p> <p>Use of $\Delta E = c^2 \Delta m$ (1)</p> <p>Use of $1 \text{ J} = 1.6 \times 10^{-19} \text{ eV}$ (1)</p> <p>$1 \text{ u} = 934 \text{ (MeV)}$ (1)</p> <p><u>Example of calculation</u></p> $\Delta E = (3.0 \times 10^8 \text{ m s}^{-1})^2 \times 1.66 \times 10^{-27} \text{ kg} = 1.494 \times 10^{-10} \text{ J}$ $\therefore \Delta E = \frac{1.494 \times 10^{-10} \text{ J}}{1.6 \times 10^{-13} \text{ J MeV}^{-1}} = 934 \text{ MeV}$	4
20(c)	<p>Use of $1 \text{ u} = 934 \text{ MeV}$ (ecf from (b) [Accept calculation from first principles] (1)</p> <p>The mass of the Fr nucleus is much greater than the mass of the α (1)</p> <p>Momentum is conserved so (recoil) velocity of Fr nucleus is much less than the velocity of the α (1)</p> <p>So the kinetic energy of the α is much greater than the kinetic energy of the Fr Or (after the decay) the α has most of the kinetic energy (1) [MP4 dependent upon MP2 or MP3]</p> <p>OR</p> <p>Use of $1 \text{ u} = 934 \text{ MeV}$ (ecf from (b) [Accept calculation from first principles] (1)</p> <p>Mathematical statement of momentum conservation (1)</p> <p>Use of $E_k = \frac{p^2}{2m}$ (1) Or use of $E_k = \frac{1}{2}mv^2$ and $p = mv$ (1)</p> <p>E_k calculated and statement that E_k is just less than 5.9 MeV (1) Or E_k calculated and statement that α has most of the kinetic energy (1)</p> <p><u>Example of calculation</u></p> $\Delta E = 6.35 \times 10^{-3} \text{ u} \times 934 \text{ MeV u}^{-1} = 5.93 \text{ MeV}$ <p>[5.91 MeV if “show that” value used]</p>	4

20(d)	Use of $\lambda t_{1/2} = \ln 2$	(1)	4
	Use of $A = -\lambda N$	(1)	
	Use of $N = N_0 e^{-\lambda t}$	(1)	
	$N = 5.6 \times 10^{13}$	(1)	
	<u>Example of calculation</u>		
	$\lambda = \frac{\ln 2}{9.9 \times 24 \times 3600 \text{ s}} = 8.10 \times 10^{-7} \text{ s}^{-1}$		
	$N = \frac{7.4 \times 10^7 \text{ s}^{-1}}{8.10 \times 10^{-7} \text{ s}^{-1}} = 9.13 \times 10^{13}$		
	$N = 9.13 \times 10^{13} \times e^{-8.10 \times 10^{-7} \text{ s}^{-1} \times 7.0 \times 24 \times 3600 \text{ s}} = 5.59 \times 10^{13}$		
Total for question 20			14