



Pearson  
Edexcel

## Mark Scheme (Results)

Summer 2022

Pearson Edexcel International Advanced Level  
In Statistics S1 (WST01) Paper 01

Question Number	Scheme		Marks
1(a)	$w = 8$		B1
	$x = 19$		B1
	$y = 37$		B1
			(3)
(b)	$"37" + 1 \times ("37" - "19") [= 55]$		M1
	59 and 64		A1ft
			(2)
(c)			M1 A1ft A1ft
			(3)
(d)	$\frac{("37" - 26) - (26 - "19")}{("37" - "19")}$		M1
	$= 0.22$ (to 2 sf)		A1
			(2)
(e)	E.g. 'The mean uses all the data'		B1
			(1)
			<b>Total 11</b>
<b>Notes</b>			
(a)	<b>B1B1B1</b>	Caution May be seen in table before part (a). $w = 28$ is first B0.	
(b)	<b>M1</b>	Calculation for the outliers using their lower quartile and upper quartile. Allow "their upper quartile" + "their IQR" for this mark i.e. $37 + 18$	
	<b>A1ft</b>	For identifying 59 and 64 as outliers from correct working. For the identification of outlier(s) (if any) from "their 55" from their shown calculation. Answer only is M0A0.	
(c)	<b>M1</b>	For a box with at least one whisker drawn	
	<b>A1ft</b>	14 for lowest whisker, 26 for median, "19" and "37" plotted for quartiles for their values for quartiles	
	<b>A1ft</b>	Upper whisker at 51 or "their 51" <b>plus</b> "their outliers" plotted but there must be at least one outlier for this mark. Condone upper whisker at "their 55". NB award A0 if there is more than one whisker at either end	
(d)	<b>M1</b>	For substituting their values into the formula	
	<b>A1</b>	Allow awrt 0.22 (allow $\frac{2}{9}$ or 0.2)	
(e)	<b>B1</b>	A correct reason which supports Landacre's use of the mean or rejects their use of the median. Allow comment relating to (slight) positive skew so mean > median so Landacre will know the larger average they will have to pay. Comments about skewness/symmetry on their own score B0. Mean includes the outliers is B1. Condone Median is not affected by the outliers for B1. Mean is more accurate is B0.	

Question Number	Scheme		Marks
2(a)	$S_{gg} = 3624.41 - \frac{144.84^2}{9} [= 1293.4516]$		M1
	$r = \frac{40.25}{\sqrt{1293.4516 \times 1.29}}$		M1
	= 0.985....		awrt 0.985 A1
			(3)
(b)	As the population/ $t$ increases, GDP/ $g$ increases oe		B1
			(1)
(c)	$b = \frac{40.25}{1.29} [= 31.20155...]$		M1
	$a = \frac{144.84}{9} - 31.20155... \times \frac{7.87}{9} [= -11.19068....]$		M1
	$g = ...31.20155...t$		A1
	$g = -11.2 + 31.2t$		A1
			(4)
(d)	The <b>GDP/<math>g</math></b> increases by (an average of) " <b>31.2</b> " billion [dollars] when the <b>population/<math>t</math></b> increases by <b>one million</b> .		B1
			(1)
(e)(i)	"-11.2" + "31.2" × 7		M1
	= 207.2...		awrt 207 A1
			(2)
(ii)	Unreliable as 7 000 000 is much greater than the mean population/ $\bar{t}$ for the 9 years.		B1
			(1)
(f)	$0.1 = 31.2x$		M1
	$x = 0.003205... \text{ million people}$		awrt 0.0032 A1
			(2)
<b>Notes</b>			<b>Total 14</b>
(a)	<b>M1</b>	Correct method for finding $S_{gg}$ (implied by awrt 1290 to 3sf)	
	<b>M1</b>	Correct method for finding $r$ using their $S_{gg}$ If $S_{gg} = 3624.41$ is used here, then M0.	
	<b>A1</b>	awrt 0.985 (correct answer only scores M1M1A1)	
(b)	<b>B1</b>	A correct interpreted contextual statement including population (or $t$ ) and GDP (or $g$ ). 'Strong positive correlation between population and GDP' on its own is B0.	
(c)	<b>M1</b>	Correct method for finding $b$	
	<b>M1</b>	Correct method for finding $a$ using their $b$ $a = 16.0... - 31.20155... \times 0.874... [= -11.19068....]$	
	<b>A1</b>	Only dep on 1 <sup>st</sup> M1 awrt 31.2 in a regression equation (allow any variables in the equation).	
	<b>A1</b>	Correct equation $g = \text{awrt } -11.2 + \text{awrt } 31.2t$ (must be $g$ and $t$ ) Do not allow fractions.	
(d)	<b>B1</b>	Idea that the GDP increases by "Their $b$ " billion dollars for every 1 million increase in population	
(e)(i)	<b>M1</b>	Correct method. Allow substitution of 7 000 000 instead of 7	
	<b>A1</b>	awrt 207 (billion) (isw after an answer of 207 is seen)	
(ii)	<b>B1</b>	Unreliable with a correct supporting comment which must reference $t$ or $\bar{t}$ [=0.874] or population 'Extrapolation so unreliable' on its own is B0. Reference to $g$ out of range is also B0.	
(f)	<b>M1</b>	Equating 0.1 with "their $b$ " × $x$ Or substituting two values for $g$ with a difference of 0.1 in their equation leading to a value of $x$	
	<b>A1</b>	awrt 0.0032 (million) Allow awrt 3200 (to 2sf) Do not allow fractions.	

Qu. No.	Scheme		Marks
3(a)	Width = 2.5 (cm)		B1
	Height = $\left(\frac{35}{15} \times 6\right) \div "2.5"$ or $\frac{4}{5} \times 7$ or $6\text{cm}^2 = 15$ (logs) or $14\text{cm}^2 = 35$ (logs) oe		M1
	= 5.6 (cm)		A1
			(3)
(b)	Median = $[20] + \frac{25}{35} \times 5$ allow $[20] + \frac{25.5}{35} \times 5$		M1
	= 23.57... allow 23.64.... awrt 23.6		A1
			(2)
(c)	19 + 35 + 8 (= 62*) or 88 - 3 - 15 - 8 (= 62*)		B1cso*
			(1)
(d)	P(4 logs fit) = $\left(\frac{62}{88}\right) \times \left(\frac{61}{87}\right) \times \left(\frac{60}{86}\right) \times \left(\frac{59}{85}\right)$		M1
	= 0.23922.... awrt 0.239		A1
			(2)
(e)(i)	mean of $y = \frac{924}{88} [= 10.5]$		M1
	mean of $w = ("10.5" + 255) \times 2$		M1
	= 531		A1
			(3)
(ii)	variance of $y = \frac{12862}{88} - ("10.5")^2 [= 35.909...]$		M1
	variance of $w = "35.909" \times 4$ or $"35.909" \div 0.5^2$		M1
	= 143.636... awrt 144		A1
			(3)
<b>Notes</b>			<b>Total 14</b>
(a)	<b>B1</b>	2.5 oe	
	<b>M1</b>	Correct method to relate area to number of logs (may be implied by "their w" $\times$ "their h" = 14)	
	<b>A1</b>	5.6 oe	
(b)	<b>M1</b>	For a correct fraction multiplied by 5. Allow working downwards e.g. $[25] - \frac{10}{35} \times 5$	
	<b>A1</b>	Correct answer from correct working. Allow exact fraction.	
(c)	<b>B1</b>	A correct calculation seen. Allow $\frac{1}{2}(16)$ for 8. Allow equivalent methods $\frac{x-54}{70-54} = \frac{26-25}{27-25}$ Minimum working required $54 + 8$ or $70 - 8$ or $\frac{54+70}{2}$ NB: $26 + x = 88 \rightarrow x = 62$ is B0.	
(d)	<b>M1</b>	For $\left(\frac{n}{88}\right) \times \left(\frac{n-1}{87}\right) \times \left(\frac{n-2}{86}\right) \times \left(\frac{n-3}{85}\right)$ (allow any $n < 88$ )	
	<b>A1</b>	awrt 0.239	
	<b>SC</b>	With replacement awrt 0.246 scores M1A0	
(e)(i)	<b>M1</b>	Correct method for finding mean of $y$ (implied by 10.5) or for equation $0.5 \sum w - 88 \times 255 = 924$	
	<b>M1</b>	Correct method for finding mean of $w$ or $\sum w = 46728$ and $\frac{46728}{88}$	
	<b>A1</b>	Cao	
(ii)	<b>M1</b>	Correct method for finding variance of $y$ (implied by awrt 35.9) or $0.25 \sum w^2 + 255^2 \times 88 - 255 \sum w = 12862$	
	<b>M1</b>	Correct method for finding variance of $w$ or $\sum w^2 = 24825208$ and $\frac{24825208}{88} - "531"'^2$	

Question Number	Scheme		Marks
	<b>A1</b>	awrt 144	
4(a)	$P(H \cup W) = P(H) + P(W) - P(H \cap W)$	$P(H' \cap W) = P(H \cup W) - P(H)$	M1
	$P(H \cap W) = \frac{3}{8} \times P(W)$	$P(H' \cap W) = P(H') \times P(W)$	M1
	$\frac{3}{4} = \frac{3}{8} + P(W) - \frac{3}{8} P(W)$	$\frac{3}{8} = \frac{5}{8} P(W)$	A1
	$P(W) = \frac{3}{5}^*$	$P(W) = \frac{3}{5}^*$	A1cso*
			<b>(4)</b>
(b)	$P(N'   H) = \frac{\frac{3}{8} - \frac{1}{15}}{\frac{3}{8}}$ or $\frac{\frac{9}{40} + \frac{1}{12}}{\frac{3}{8}}$ or $1 - \frac{1}{15}$		M1
	$= \frac{37}{45}$ = awrt 0.822		A1
			<b>(2)</b>
(c)			B1 M1 M1 M1 A1 <b>(5)</b>
<b>Notes</b>			<b>Total 11</b>
(a)	<b>M1</b>	for use of $P(H \cup W) = P(H) + P(W) - P(H \cap W)$ (with at least one value correctly substituted) or use of $P(H' \cap W) = P(H \cup W) - P(H)$ (with at least one value correctly substituted)	
	<b>M1</b>	for use of $P(H \cap W) = P(H) \times P(W)$ or use of $P(H' \cap W) = P(H') \times P(W)$	
	<b>A1</b>	a correct equation in $P(W)$ (allow $W$ or $x$ here)	
	<b>A1cso*</b>	Correct solution ending with $P(W) = \frac{3}{5}$ with no wrong working seen. Dep. on all previous marks.	
	<b>NB</b>	A method which uses $\frac{3}{5}$ or $\frac{3}{5} \times \frac{3}{8} \left[ = \frac{9}{40} \right]$ can score maximum M1M1A0A0.	
(b)	<b>M1</b>	For $\frac{p}{\frac{3}{8}}$ where $0 < p < \frac{3}{8}$ use of independence is M0 e.g. $\frac{x \times \frac{3}{8}}{\frac{3}{8}}$	
	<b>A1</b>	awrt 0.822	
(c)	<b>B1</b>	3 circles labelled. Either $N$ inside $H$ or allow as intersecting circles with $P(N \cap H') = 0$ , but do not allow blank space to be considered 0. Condone missing box for this mark. Allow all 3 circles overlapping with all zeros correctly labelled.	
	<b>M1</b>	For $P(H \cap W) = \frac{3}{8} \times \frac{3}{5} \left[ = \frac{9}{40} \right]$ seen or correctly placed in Venn diagram.	
	<b>M1</b>	For their $\frac{3}{5} - \frac{9}{40} = \frac{3}{8}$ (may be implied by the regions in their $P(W)$ adding to 0.6)	
	<b>M1</b>	For their $\frac{3}{8} - \frac{9}{40} - \frac{1}{15} = \frac{1}{12}$	
	<b>A1</b>	Fully correct diagram with $\frac{1}{4}$ and box and correct probabilities (allow exact decimal equivalents)	

Question Number	Scheme		Marks
5(a)	$[E(R^2)] = 2^2 \times 0.25 + 3^2 \times 0.3 + 4^2 \times 0.15 + 5^2 \times 0.1 + 6^2 \times 0.2 (= 15.8^*)$		B1cso* (1)
(b)	$[sd(R)] = \sqrt{15.8 - 3.7^2}$		M1
	$= \sqrt{2.11}$		
	Standard deviation = 1.4525....	awrt 1.45	A1 (2)
(c)	$d = 1$		B1 (1)
(d)	$0.1 + 0.2 + 0.1 + a + b = 1$ oe		M1
	$2 \times 0.1 + 3 \times 0.2 + 4 \times 0.1 + 5a + 6b = 4.55$ oe		M1
	$5(0.6 - b) + 6b = 3.35$ or $5a + 6(0.6 - a) = 3.35 \Rightarrow a = 0.25$ or $b = 0.35$		M1
	$c = 0.4 + "0.25"$ or $c = 1 - "0.35"$		M1
	$c = 0.65$ oe		A1 (5)
(e)	$0.9 \times 0.75 \times 0.1$		M1
	$= 0.0675$		A1
			(2)
(f)	For identifying that if Jessie scores 2, Pabel has no spin oe may be implied		M1
	$[0.10 \times 0 + ] 0.2 \times 0.3 + 0.1 \times 0.15 + "0.25" \times 0.1 + "0.35" \times 0.2$		M1
	$= 0.17$		A1
			(3)
<b>Notes</b>			<b>Total 14</b>
(a)	<b>B1</b>	Correct calculation with all products seen (allow $1 + 2.7 + 2.4 + 2.5 + 7.2$ ) Figures may be seen in table before part (a). Condone missing addition signs if products seen in table.	
(b)	<b>M1</b>	Use of formula including the square root	
	<b>A1</b>	awrt 1.45 (correct answer with no working scores M1A1)	
(c)	<b>B1</b>	For 1	
(d)	<b>M1</b>	Allow equivalents eg $a + b = 0.6$	
	<b>M1</b>	Allow equivalents eg $5a + 6b = 3.35$	
	<b>M1</b>	Correct method to eliminate $a$ or $b$ (implied by a correct value for $a$ or $b$ ) This mark can still be scored even if the method leads to a value of $a$ or $b$ which is not a probability. May see $a = c - 0.4$ to eliminate $a$ or $b = 1 - c$ used to eliminate $b$	
	<b>M1</b>	A complete method for finding the value of $c$ (condone using any value of $a$ and $b$ for this mark)	
	<b>A1</b>	0.65 oe	
(e)	<b>M1</b>	For the product of 3 probabilities	
	<b>A1</b>	0.0675 or exact equivalent fraction eg $\frac{27}{400}$	
(f)	<b>M1</b>	Identifying that if Jessie scores 2, there is only one spin or the 4 correct possibilities only	
	<b>M1</b>	At least 3 correct non-zero probability products ft their $a$ and $b$ (an answer of 0.195 scores M0M1A0)	
	<b>A1</b>	0.17	

Question Number	Scheme		Marks
6(a)	$P(V > 104.9) = P\left(Z > \frac{104.9 - 100}{2.5}\right)$		M1
	$= 1 - 0.975$		M1
	$= 0.0250$	0.025 or awrt 0.0250	A1
			(3)
(b)	Expected number = $150 \times "0.025"$		M1
	$= 3.75$	awrt 3.75	A1
			(2)
(c)	$[P(V > v   V < 104.9)] = \frac{P(v < V < 104.9)}{P(V < 104.9)} = 0.2801$		M1
	$P(v < V < 104.9) = (1 - "0.025") - P(V < v)$		M1
	$P(V < v) = (1 - "0.025") - (1 - "0.025") \times 0.2801 [=0.7019] \text{ oe}$		dM1
	$\frac{v - 100}{2.5} = 0.53$		M1A1
	$v = 101.325$	awrt 101.32 or awrt 101.33	A1
		(6)	
<b>Notes</b>			<b>Total 11</b>
(a)	<b>M1</b>	Standardising with 104.9, 100 and 2.5 (allow $\pm$ ) implied by 1.96 seen	
	<b>M1</b>	For use of $1 - p$ with $0.9 < p < 1$ condone answer of 0.0249 for this mark	
	<b>A1</b>	Allow 0.025 or awrt 0.0250 (NB calculator answer is 0.02499...) (answer only scores M1M1A1)	
(b)	<b>M1</b>	For $150 \times$ "their part (a)"	
	<b>A1</b>	awrt 3.75 isw after answer of 3.75 seen. If 3.75 not seen, allow 4 if the method mark is awarded.	
(c)	<b>M1</b>	For writing or using a ratio of probabilities with denominator $P(V < 104.9)$ oe and equating to 0.2801 $\frac{P}{P(V < 104.9)} = 0.2801$ implied by awrt 0.273 Use of independence is M0 e.g. $\frac{x \times P(V < 104.9)}{P(V < 104.9)} = 0.2801$	
	<b>M1</b>	For writing or using $P(V > v \cap V < 104.9) = P(V < 104.9) - P(V < v) [= (1 - "0.025") - P(V < v)]$	
	<b>ALT</b>	<b>For first two M marks</b> $\frac{P(V < v)}{P(V < 104.9)} = 1 - 0.2801$ scores M1M1, then follow scheme.	
	<b>dM1</b>	(dep M1) Dependent on previous M1 for rearranging to find $P(V < v)$ Allow equivalent oe $(1 - "their(a)")(1 - 0.2801)$ NB: $[P(V < v) =]$ awrt 0.702 implies M1M1M1 or $[P(V > v) =]$ awrt 0.298 implies M1M1M1	
	<b>M1</b>	Standardising with 100, 2.5 and equating to a z-value, $\frac{v - 100}{2.5} = z$ $0.4 <  z  < 0.6$ Watch out for $\frac{v - 100}{2.5} =$ probability which is M0	
	<b>A1</b>	Correct equation with compatible signs	
	<b>A1</b>	awrt 101.33 (allow awrt 101.32 from use of calculator)	