



Mark Scheme (Final)

October 2019

Pearson Edexcel International Advanced Level
In Statistics S1 (WST01/01)

| Question Number | Scheme | Marks |
|-----------------|--|--|
| 1(a) | $[S_{xy} =] = \underline{1818}$ | B1 (1) |
| (b) | $b = \frac{1818}{1754} [=1.036 \text{ or } \frac{909}{877}]$ ALT $c = -1.036... \times \frac{5}{8}$ or $\bar{f} = 80 \times \frac{5}{8} + 3500$ $= -0.6478$ or $\bar{h} = 4500$ and $\bar{f} = 3550$ $y = 1.04x - 0.6478...$ or eqn of form $h = ... + 1.04f$ $\frac{h-4500}{80} = 1.036 \times \frac{f-3500}{80} - 0.6479...$ or $4500 - 1.04 \times 3550 = (820.467...)$ $h = 820 + 1.04f$ | M1 M1 A1 A1ft dM1 A1 (6) |
| (c) | <p>On average for every <u>increase in area of 1</u> (m²) the annual heating <u>bill increases</u> by approximately (\$) <u>1.04</u></p> <p>The cost of heating is (\$) 820 even if there is no floor space (no building)</p> <p><u>or</u> the “standing charge” / “base rate” etc is (\$) 820</p> | B1ft B1ft (2) |
| (d) | $"820.46..." + "1.036..." \times 4600$ $= (\$) 5588.31...$ ans in range (5560-5610) | M1 A1 (2) |
| Total 11 | | |

| Notes | | |
|-------|--|--|
| (b) | <p>1st M1 for use of $\frac{S_{xy}}{S_{xx}}$ ft their S_{xy} (allow correct use of S_{fh} etc)</p> <p>2nd M1 for $c = -(\text{their } b) \times \frac{5}{8}$ or ALT for a correct expression for \bar{f}</p> <p>1st A1 awrt -0.648 or ALT for correct values for both \bar{h} and \bar{f}</p> <p>2nd A1ft for equ'n in the form $y = c + bx$ with their c and $b =$ awrt 1.04 (dep on 1st M1 only) or ALT for $h = ... + (\text{awrt } 1.04)f$ [ignore any intercept]</p> <p>3rd dM1 (dep on 2nd M1) for substituting $\frac{h-4500}{80}$ and $\frac{f-3500}{80}$ into their regression line of y on x or ALT for correct method for intercept in h, f equation – ft their 1.04</p> <p>3rd A1 for $h =$ awrt $820 +$ awrt $1.04f$ [Do not allow fractions for this final mark]</p> | |
| (c) | Must see correct words used, not just letters f and h | |
| (i) | B1ft for correct interpretation need: “increase of area by 1” and “increase in bill by “1.04”” Can be other words giving same idea but must see their 1.04 (can ignore units) | |
| (ii) | B1ft for a suitable explanation e.g. “cost if area is zero” Must see “their 820” (which must be > 0) but can ignore units. Accept reasonable alternatives to “standing charge” | |
| (d) | M1 for substituting 4600 into their regress' line of h on f (or use of $x = 13.75$ in y, x equation) A1 allow answers in range 5580 - 5610 | |

| Question Number | Scheme | | Marks |
|-----------------|---|--|-----------------|
| 2(a) | $10 \times 7.8 \div 1.2 / 10 \times 7.8 \times \frac{5}{6} / \frac{1}{6} \times 10 \times 39$ | $5 \times 3.6 \div 1.2 / 5 \times 3.6 \times \frac{5}{6} / \frac{1}{6} \times 5 \times 18$ | M1 |
| | (Time 45-55 freq =) 65 | (Time 55-60 freq =) 15 | A1 |
| | $260 - "65 + 180" = \underline{15}$ | $260 - "15 + 180" = \underline{65}$ | B1ft (3) |
| (b)(i) | $\left[\text{Mean} = \frac{11087.5}{260} \right] = 42.644\dots$ | awrt 42.6 | B1 (1) |
| (ii) | Standard deviation = $\sqrt{\frac{505718.75}{260} - \left(\frac{11087.5}{260}\right)^2} = 11.249\dots (s = 11.27\dots)$ | awrt 11.2 | M1 A1 (2) |
| (c) | Median = $(30) + 15 \times \frac{130 - 20}{145}$ or $(45) - 15 \times \frac{(165 - 130)}{145}$ | $= 41.379\dots$ awrt 41.4 | M1 A1 (2) |
| (d) | Positive (skew) since the median is less than the mean or Positive (skew) since the histogram has a "tail" on the right oe (Allow use of quartiles if $Q_1 = [34.65\dots]$ and $Q_3 = [49.61\dots]$ are correct to 2sf) | | B1 (1) |
| (e) | Number of people = $\frac{6}{15} \times 145 + 20$ [= 78] | | M1 |
| | Probability being less than 36 seconds = $\frac{78}{260} = 0.3$ o.e. | | A1 |
| | $\left(\frac{"78"}{260}\right) \times \left(\frac{"77"}{259}\right) \times \left(\frac{"76"}{258}\right); = \frac{209}{7955}$ or 0.02627... | awrt 0.0263 | M1;A1 (4) |
| Total 13 | | | |
| Notes | | | |
| (a) | M1 for a correct expression for one frequency A1 for 65 or 15 B1 for 15 or 65 ft <u>one of</u> their incorrect frequency or their "180" | Look in table in QP If table and text disagree take table. | |
| (b)(i) | B1 for awrt 42.6 [Beware use of units e.g. 42.6s which looks like 42.65] (Allow $\frac{4435}{104}$) | | |
| (ii) | M1 ft their mean. Must have the square root. [Allow use of their $\Sigma ft^2 > 300\ 000$ here] A1 for awrt 11.2 [but allow 11.25] (allow $s = 11.3$) [Answer only 2/2] | | |
| NB | [Beware correct formula with mean = 42.64 gives 11.265 = 11.3 (3sf) but scores M1A0] | | |
| (c) | M1 for correct use of linear interpolation to find the median ignore the end points (may be implied by correct answer). NB may work up or down and allow use of 130.5 A1 for awrt 41.4 (Use of 130.5 gives 41.43...) [Again beware of units 41.4s etc] | | |
| (d) | B1 positive (skew) with either reason [No ft for -ve skew if $Q_2 > \text{mean}$]. Ignore use of mode and condone "positive correlation" If $ Q_2 - \bar{t} < 1$ allow mean close to median so no skew or symmetric. | | |
| (e) | 1 st M1 for a correct method for finding the number of people. [Correct expression or 78] 1 st A1 for the correct probability of 0.3 or exact equivalent 2 nd M1 for use of <u>their</u> probability without replacement (condone with replacement) Score this mark for p^3 where $p =$ their 0.3 from <u>any</u> method e.g. use of normal etc 2 nd A1 for awrt 0.0263 | | |

| Question Number | Scheme | Marks |
|-----------------|---|---|
| 3(a) | $P(X < 40) = P\left(Z < \frac{40 - 42}{5}\right) = P(Z < -0.4)$ $= 1 - 0.6554$ $= 0.3446$ | M1 M1 A1 (3) |
| (b) | <p>P(Qualify) = $1 - ("0.3446")^3$ <u>or</u> $(1 - "0.3446") + ("0.3446")(1 - "0.3446") + ("0.3446")^2(1 - "0.3446")$ [$q = 0.9590\dots$ full calc: 0.9590867...]</p> $P(X > 45) = P\left(Z > \frac{45 - 42}{5}\right) = P(Z > 0.6)$ $= 1 - 0.7257$ <u>or</u> 0.2743 allow 1 - awrt 0.726 <u>or</u> awrt 0.274 $P(X > 45 \text{ on 3rd throw} \mid \text{in final}) = \frac{"0.3446" \times "0.2743"}{"0.959"}$ $= \frac{"0.0326\dots"}{"0.9590\dots"} \quad (\text{calc: } 0.033952\dots)$ | M1 M1 A1 dM1 A1 (5) Total 8 |
| Notes | | |
| (a) | 1 st M1 for standardising with 40, 42 and 5. Allow $\pm \frac{40 - 42}{5}$ 2 nd M1 for $1 - p$ (where $p > 0.5$) A1 awrt 0.345 (NB Calc gives 0.3445783...) | |
| (b) | 1 st M1 for identifying <u>all</u> the cases to qualify with correct ft probabilities 2 nd M1 for an attempt standardise with 45, 42 and 5 ie $\pm \frac{45 - 42}{5}$ SC B1 Use $\mu = 40$ Using $\mu = 40$ to find $P(X > 45)$ will give 0.1587 and can award B1 if we see this <u>used</u> in an expression of the form $(\text{"0.3446"})^2 \times 0.1587$. Score on open as 2 nd M0 1 st A1 They may also be able to score 1 st M1 and 3 rd M1 as well 1 st A1 for 1 - awrt 0.726 or awrt 0.274 (sight of either of these scores 2 nd M1 and 1 st A1) May be part of an expression such as $(\text{"0.3446"})^2 \times 0.2743$ 3 rd dM1 dep on 1st M1 for $\frac{(\text{"their 0.3446"})^2 \times \text{"their 0.2743"}}{\text{their } q}$ providing the numerator < denominator and num and denom are both probs 2 nd A1 for awrt 0.034 (NB numerator is awrt 0.033 so <u>must</u> be awrt 0.034) | |

| Question Number | Scheme | Marks |
|-----------------|--|------------------------------------|
| 4(a) | <p style="text-align: center;">If any part, especially (a) or (b), is missing send to review</p> 0.72 | B1 (1) |
| (b) | C (is most likely to be the 100 metre junior champion) | B1 (1) |
| (c) (i) | $S_{xx} = 3445.26 - \frac{164.4^2}{8} \left[= 66.84 \text{ or } \frac{1671}{25} \right]$ $r = \frac{60.85}{\sqrt{66.84 \times 67.52}}$ $= 0.90578\dots \qquad \text{awrt } \underline{\mathbf{0.906}}$ | M1 M1 A1 (3) |
| (ii) | The faster boys are in the the 100 metres, the faster they are in the 200 metres | B1 (1) Total 6 |
| Notes | | |
| (c) (i) | <p>1st M1 for a correct expression, allow the use of $n = 10$, ie $S_{xx} = 3445.26 - \frac{164.4^2}{10} [= 742.524]$</p> <p>Condone one slip e.g. 3445.6 instead of 3445.26 etc</p> <p>2nd M1 for an attempt at a correct formula for r using S_{yy} and S_{xy} and their S_{xx}</p> <p>Condone one slip e.g. 60.84 or 66.48 miscopied for 66.84</p> <p>A1 for awrt 0.906</p> <p>NB Use of $S_{xx} = 742.524$ gives $r = 0.272$ and can score M1M1A0 provided expressions are seen for S_{xx} and r</p> | |
| (ii) | <p>NB on open this is an A1 mark but we are treating it as a B1 It does not depend on M1 in (c)(i)</p> <p>B1 allow equivalent statements e.g. on average boys that are faster/slower in the 100 metres are also faster/slower in the 200 metres</p> <p>Comment must be: (1) a comparison of <u>time</u> e.g. faster, quicker, slower etc (not “higher”) and (2) mention <u>100 metres</u> or <u>200 metres</u> (and imply the other)</p> | |

| Question Number | Scheme | Marks |
|-----------------|--|---|
| 5(a)(i) | $P(D) = \frac{200}{320} = \frac{5}{8}$ (or exact equivalent e.g. 0.625) | B1 (1) |
| (ii) | $P(D \cap X') = \frac{1}{2}$ oe | B1 (1) |
| (iii) | $P(D' \cup Z') = \frac{320-88}{320}; = \frac{29}{40} = 0.725$ o.e. | M1; A1 (2) |
| (b) | $P(Z D) = \frac{\frac{88}{320}}{\frac{200}{320}}; = \frac{88}{200}$ or $\frac{11}{25}$ or 0.44 oe | M1; A1 (2) |
| (c) | X and Y or X and Z or Y and Z (Allow X, Y etc) | B1 (1) |
| (d) | $P(D) \times P(X) = 0.625 \times 0.2$ or $\frac{5}{8} \times \frac{64}{320} = 0.125 = P(D \cap X)$ or $P(D X) = \frac{40}{64} = 0.625 = P(D)$ or $P(X D) = \frac{40}{200} = \frac{1}{5} = P(X) = \frac{24+40}{320}$ So yes they are independent | M1 A1 (2) |
| (e)(i) | A house that does not have a driveway but has exactly two cars | B1 (1) |
| (ii) | A house that has a driveway (with) fewer than two cars (oe) | B1 B1 (2) Total 12 |
| Notes | | |
| (a)(iii) | <p style="text-align: center;">If any part(s) of this question are missing please send to review</p> <p>M1 for identifying the correct 7 values: 24, 40, 35, 37, 32, 44 and 20 or sum of 232 A1 for $\frac{29}{40}$ or exact equivalent e.g. 0.725</p> <p>(b) M1 for a ratio of probabilities with numerator of $\frac{88}{320}$ and denominator of their (a)(i) A1 for 0.44 or exact equivalent</p> <p>(c) B1 for at least one correct pair and no incorrect ones. Do not allow e.g. $P(Y \cap Z)[=0]$ etc</p> <p>(d) M1 for a correct test with all required probs (labels and values) stated or implied - ft $P(D)$ A1 for a correct conclusion – allow “yes they are” but must be events not probabilities e.g. a conclusion that $P(D)$ and $P(X)$ are independent is A0</p> <p>(e)(ii) 1st B1 for a house that has a driveway 2nd B1 for fewer than two cars (Allow 0 or 1 but must not include both no car and 1 car) e.g. “has a driveway with 1 car <u>and</u> has a driveway with no car” is B1B0 but “has a driveway with 1 car <u>or</u> has a driveway with no car” is B1B1</p> | |

| Question Number | Scheme | Marks |
|---------------------|---|--|
| <p>6 (a)</p> | $\frac{3.968 - \mu}{\sigma} = -1.2816 \quad \text{or} \quad \frac{4.026 - \mu}{\sigma} = 1.0364$ $\mu - 1.2816\sigma = 3.968 \quad (\text{Calc: } -1.28155156\dots)$ $\mu + 1.0364\sigma = 4.026 \quad (\text{Calc: } 1.03643338\dots)$ $2.318\sigma = 0.058$ $\sigma = 0.0250\dots \quad \mu = 4.00\dots \quad \text{awrt } \underline{\mathbf{0.025 \text{ and } 4}}$ | <p>M1A1A1</p> <p>dM1</p> <p>A1</p> <p>(5)</p> |
| <p>(b)</p> | $Q_3 = \text{awrt } 30.3 \text{ (calc: } 30.337\dots) \text{ or } Q_3 - Q_1 = \text{awrt } 0.6 \text{ oe (calc: } 0.6744\dots)$ $30.3 + 1.5("30.3" - 29.7)[= 31.2] \quad \text{or} \quad 29.7 - 1.5("30.3" - 29.7)[= 28.8]$ $P(L > "31.2") = P\left(Z > \frac{"31.2" - 30}{0.5}\right) \quad \text{or} \quad P(L < "28.8") = P\left(Z < \frac{"28.8" - 30}{0.5}\right)$ $= 0.0082$ <p>Probability it is an outlier = 2×0.0082</p> $= 0.0164 \quad \text{answer in range } \underline{\mathbf{(0.006 \sim 0.017)}}$ | <p>B1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(5)</p> <p>Total 10</p> |
| Notes | | |
| <p>(a)</p> | <p>1st M1 for standardising with μ and σ and forming an equation in μ and σ with $z > 1$</p> <p>1st A1 for one correct equation in any form with z value as given or better</p> <p>2nd A1 for a 2nd correct equation in any form allow 2dp or better for the z value</p> <p>2nd dM1 (dep on 1st M1) for correct method to solve* their 2 linear, simultaneous equations. Can be implied by both correct answers. [*Must see correct substitution or correct addition/subtraction of all 3 terms]</p> <p>2nd A1 for both $\mu = \text{awrt } 4$ and $\sigma = \text{awrt } 0.025$ [Check it follows from their working.] NB Could score M1A0A1M1A1 or M1A1A0M1A1 here</p> | |
| <p>(b)</p> | <p>B1 awrt 30.3 or IQR = awrt 0.6 or awrt 0.67</p> <p>1st M1 correct method for finding 1 outlier limit – fit their Q_3 or their IQR</p> <p>2nd M1 standardising with their limit, 30 and 0.5 allow \pm leading to a probability < 0.05 Can be implied by a correct probability statement e.g. $P(L < 28.8) = 0.0082$</p> <p>3rd M1 multiplying their probability by 2 (or adding their two probs both < 0.05)</p> <p>A1 (dependent on all 3 M marks) for an answer in the range $0.006 \sim 0.017$</p> | |
| <p>Calc</p> | <p>Use of full calc values: If they use a calculator the lower limit is 28.651... upper limit is 31.3489 and probability comes to $2 \times 0.00348835\dots = \text{awrt } 0.00698$</p> | |

| Question Number | Scheme | Marks | | | | | | | | | | | | | | | | | | |
|-----------------|---|------------------------------|----------------|-----------------|----------------|---|---|-----|----|-----|-----|-----|-----|------------|----------------|----------------|----------------|-----------------|----------------|--------------------|
| 7(a)(i) | $2a + 2b = 0.5$ oe ; $5a + 11b = 1.55$ oe (any unsimplified form) e.g. $5a + 11(0.25 - a) = 1.55$ [implies $6a = 1.2$ oe] $a = 0.2^*$ | B1; B1 M1 A1 cso B1 | | | | | | | | | | | | | | | | | | |
| (ii) | $b = \underline{0.05}$ | (5) | | | | | | | | | | | | | | | | | | |
| (b) | $[E(X^2) =]1^2 \times 0.35 + 2^2 \times 0.2 + 3^2 \times 0.2 + 4^2 \times 0.15 + 5^2 \times 0.05 + 6^2 \times 0.05 [=8.4]$ $\text{Var}(X) = "8.4" - 2.5^2 = [2.15]$ $\text{Var}(4X + 3) = 16 \text{Var}(X)$ $= \underline{34.4}$ | M1 M1 M1 A1 | | | | | | | | | | | | | | | | | | |
| (c) | Expected profit = 2.5×60 or $2.5 \times 80 - 2.5 \times 20$ oe = <u>150</u> (cents) or \$1.50 per customer | M1 A1 | | | | | | | | | | | | | | | | | | |
| (d) | Let W be the profit, in cents per customer <table border="1" data-bbox="280 790 1289 931"> <tr> <td>y</td> <td>1</td> <td>2</td> <td>3</td> <td>5</td> <td>6</td> </tr> <tr> <td>w</td> <td>60</td> <td>120</td> <td>180</td> <td>220</td> <td>280</td> </tr> <tr> <td>$P(Y = y)$</td> <td>$\frac{3}{40}$</td> <td>$\frac{4}{40}$</td> <td>$\frac{3}{40}$</td> <td>$\frac{22}{40}$</td> <td>$\frac{8}{40}$</td> </tr> </table> $[E(W)] = \frac{1}{40}(60 \times 3 + 120 \times 4 + 180 \times 3 + 220 \times 22 + 280 \times 8)$ $= \underline{207}$ cents per customer [May work in dollars e.g. \$2.67 or \$2.27 scores B1B0M1A0 and \$2.07 4/4] | y | 1 | 2 | 3 | 5 | 6 | w | 60 | 120 | 180 | 220 | 280 | $P(Y = y)$ | $\frac{3}{40}$ | $\frac{4}{40}$ | $\frac{3}{40}$ | $\frac{22}{40}$ | $\frac{8}{40}$ | B1; B1 M1 A1 |
| y | 1 | 2 | 3 | 5 | 6 | | | | | | | | | | | | | | | |
| w | 60 | 120 | 180 | 220 | 280 | | | | | | | | | | | | | | | |
| $P(Y = y)$ | $\frac{3}{40}$ | $\frac{4}{40}$ | $\frac{3}{40}$ | $\frac{22}{40}$ | $\frac{8}{40}$ | | | | | | | | | | | | | | | |
| Notes | | | | | | | | | | | | | | | | | | | | |
| | Mark (a)(i) and (a)(ii) together | | | | | | | | | | | | | | | | | | | |
| (a)(i) | 1 st B1 for a correct equation using $\Sigma \text{prob} = 1$ 2 nd B1 for a correct equation using $E(X) = 2.5$ M1 dep on at least B1 for <u>eliminating</u> a or b leading to a linear equation in a or b only A1cso for $a = 0.2$ <u>correctly</u> shown. Dependent on M1 scored and two correct equations seen. | | | | | | | | | | | | | | | | | | | |
| (ii) | B1 for $b = 0.05$ (or exact equivalent) Independent of other marks in (a)(i). Look by table | | | | | | | | | | | | | | | | | | | |
| (b) | 1 st M1 for an attempt at $E(X^2)$ with at least 3 correct products. Allow ft of their value of b Allow expression even if labelled $\text{Var}(X)$ but label of $\text{Var}(X)$ loses 2 nd M1 but can get 3 rd M1 2 nd M1 for use of $E(X^2) - [E(X)]^2$ ft their value of $E(X^2)$ 3 rd M1 for seeing $16 \text{Var}(X)$ [Allow this mark if clearly stated $\text{Var}(X) = E(X^2) = 8.4$] A1 for 34.4 or exact equivalent e.g. $\frac{172}{5}$ | | | | | | | | | | | | | | | | | | | |
| (c) | M1 for 2.5×60 or any other fully correct equivalent expression (allow 2.5×0.6) A1 for 150 or accept \$1.5 (working with dollars requires units) | | | | | | | | | | | | | | | | | | | |
| (d) | 1 st B1 for 1 st 3 values of W (can allow in an expression for $E(W)$) Look by table but must 2 nd B1 for last 2 values for W be in part (d) M1 for attempt at $E(W)$ ft their W values but at least 3 correct ft products Dependent on at least one of the first two B1 marks A1 for 207 | | | | | | | | | | | | | | | | | | | |
| ALT | Use of $E(Y)$ [= 4.45] with at least 3 correct products seen and $E(W) = pE(Y) - q$ 1 st B1 for $p = 60$ and 2 nd B1 for $q = \left(\frac{22}{40} + \frac{8}{40}\right) \times 80$ (or 60) M1 for $E(W)$ expression of the form above and dep on at least one B mark scored | | | | | | | | | | | | | | | | | | | |
| Total 15 | | | | | | | | | | | | | | | | | | | | |