





Mark Scheme (Results)


October 2021

Pearson Edexcel International A Level
In Statistics S1 (WST01) Paper 01

Question Number	Scheme	Marks
1 (a)	[Sum of probs = 1 gives $p + q = 0.2$ and] so $P(B) = \underline{0.5}$	B1 (1)
(b)	e.g. $P(A) = 0.3$ or $0.1 +$ "their value for $p + q$ ", $P(A \cap B) = 0.2$ or "their value for $p + q$ ", and $[P(A) \times P(B) =] 0.3 \times "0.5" [= "0.15"]$ $0.15 \neq 0.2$ so $[A$ and B are] not independent	M1 A1 (2)
(c)	$[P(C B) =] \frac{p}{"0.5"} = p + 0.06$ (o.e.) $[2p = p + 0.06$ so] $\underline{p = 0.06}$ [Use of $p + q = 0.2$ gives] $\underline{q = 0.14}$	M1 A1 A1 (3)
(d)	Suitable event D drawn. [See Venn diagrams below]	B1 (1)
Notes		
(a)	B1 for 0.5 or exact equivalent	
(b)	M1 for sight of correct probabilities for $P(A)$ and $P(A \cap B)$ clearly labelled , $0.3 \times "0.5"$ seen or $P(A) \times P(B) = 0.15$ Allow $0.04 + 0.06 + 0.2$ for $P(A)$ if clearly labelled $P(A \cap B)$ may be stated in part (a) $P(B)$ can fit from (a) eg $P(A) = "0.5" - 0.3 - 0.2$ May see $P(B A) = \frac{2}{3}$ and compared with $P(B)$ or $P(A B) = 0.4$ and $P(A) = 0.3$	
ALT	A1 For all the correct probabilities and calculations, a comparison and correct conclusion. We need to see 0.15 but will accept $P(A \cap B) \neq P(A) \times P(B)$ instead of $0.15 \neq 0.2$ for comparison SC Allow M1A0 for $P(A) = 0.1 + p + q$; $P(A \cap B) = p + q$ clearly labelled and $0.5 \times (0.1 + p + q)$ or $(p + q + 0.3)(0.1 + p + q)$ given.	
(c)	M1 fit their $P(B)$ from part (a). For a correct equation in p or q based on the given statement. NB equation in terms of q is $\frac{0.2 - q}{"0.5"} = 0.26 - q$ (o.e.) Allow $\frac{p}{0.3 + p + q} = p + 0.06$	
Ans only	1 st A1 for ($p =$) 0.06 2 nd A1 for ($q =$) 0.14 ($p =$) 0.06 and ($q =$) 0.14 3/3	
(d)	B1 for a suitable event D drawn that has an intersection with B but not with A . Condone if not labelled D	



Question Number	Scheme	Marks
<p>2 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>$[S_{xp} =] 2347 - \frac{93 \times 273}{12}$ or $2347 - \frac{25\,389}{12}$ [= 231.25] (*)</p> <p>$[S_{pp} =] 6602.72 - \frac{273^2}{12} = [391.97]$</p> <p>$[r =] \frac{231.25}{\sqrt{148.25 \times 391.97}}$ = 0.959307... awrt 0.959</p> <p>$\left[b = \frac{S_{xp}}{S_{xx}} = \right] \frac{231.25}{148.25} [= 1.559865...]$</p> <p>$a = \frac{273}{12} - 1.56 \times \frac{93}{12}$ or $22.75 - 1.56 \times 7.75$ [= 10.66...]</p> <p>$b =$ awrt 1.6 or $a =$ awrt 11</p> <p>$p = 10.7 + 1.56x$</p> <p>e.g. each extra employee costs the company (on average) "\$156" a year in paper</p> <p>[New $p =] 0.8 \times 10.66... + \frac{1.559...}{2} \times \frac{93}{12}$ [= 14.573...]</p> <p>[compared with $\bar{p} = 22.75$] so percentage saving is $\frac{22.75 - 14.573...}{22.75} [\times 100]$ = 35.94... awrt 36[%]</p>	<p>B1cso (1)</p> <p>M1</p> <p>M1</p> <p>A1 (3)</p> <p>M1</p> <p>M1</p> <p>A1 A1 (4)</p> <p>B1 (1)</p> <p>M1</p> <p>M1</p> <p>A1 (3)</p>
Notes		[12]
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>B1 for either correct expression [don't need = 231.25]</p> <p>1st M1 for attempt at correct expression for S_{pp} Allow one transcription error e.g. 6620.72 May be seen in part (a) If no correct expression seen allow $S_{pp} =$ awrt 392 or correctly placed in formula for r</p> <p>2nd M1 for a correct expression for r, ft their S_{pp}</p> <p>A1 for awrt 0.959</p> <p>1st M1 for a correct expression for b (may be implied by a correct value of awrt 1.56)</p> <p>2nd M1 for a correct expression for a ft their value for b. May be implied by awrt 10.7</p> <p>1st A1 for $b =$ awrt 1.6 or $a =$ awrt 11</p> <p>2nd A1 for correct equation in p and x with $b =$ awrt 1.56 and $a =$ awrt 10.7</p> <p>B1 for a suitable contextual comment that mentions their value of b Allow multiples eg every extra 100 employees costs the company "\$15600". Condone missing \$ sign or use of £. Do not allow "\$1.56" for 1 person unless indicates in 100's</p> <p>1st M1 for a correct expression for average value of p using new model [ft their a and b]</p> <p>2nd M1 for correct percentage saving calculation using 22.75 (e.g. $\frac{14.573...}{22.75} [\times 100]$) Allow use of $10.7 + 1.56 \times \frac{93}{12} [\approx 22.79]$ for 22.75 May be implied by correct answer.</p> <p>A1 for awrt 36</p> <p>SC use of 93 throughout part (e) rather than 7.75 leading to awrt 48 or 0.48 (they will need to use the regression line from part(c) to calculate the original value) gains M0M1A0</p> <p>SC use of 93 in part(c) Answer of 36% gains M1M1A1, 64% or 0.64 gains M1M1A0.</p>	

Question Number	Scheme	Marks
<p>3. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)(i)</p> <p>(ii)</p>	<p>[Median =] 53</p> <p>$Q_1 = 45$ $Q_3 = 61$</p> <p>[IQR =] $61 - 45 = \mathbf{16}$ (*)</p> <p>$Q_1 - 1.5 \times (\text{IQR}) = 45 - 1.5 \times 16 [= \mathbf{21}]$ or $Q_3 + 1.5 \times (\text{IQR}) = 61 + 1.5 \times 16 [= \mathbf{85}]$</p> <p>Outliers are $< \mathbf{21}$ or $> \mathbf{85}$</p> <p>So there are three outliers at 13, 87 and 88</p>  <p>e.g. the females are generally older than the men as median is higher ($67 > 53$)</p> <p>No change to box plot means one in each section so granddaughter [34~56]</p> <p>Eldest daughter in range $[67\sim 72]$ or Anja's age $[72\sim 93]$</p> <p>Since Anja 23 years older than eldest daughter Anja in range [90~93]</p>	<p>B1</p> <p>(1)</p> <p>M1</p> <p>A1cso</p> <p>(2)</p> <p>M1</p> <p>A1ft</p> <p>A1</p> <p>(3)</p> <p>M1</p> <p>A1ft</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p>B1</p> <p>(1)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>(3)</p>
Notes		[14]
(a)	B1 for 53	
(b)	M1 for an attempt at both and at least one correct. No need to be labelled.	
	A1cso for both correct quartiles seen and $61 - 45$ leading to 16	
(c)	<p>M1 for attempting at least one of the limits. Can fit their quartiles. May be implied by 85 or 21</p> <p>1st A1ft for both outlier limits correct or correct ft using their quartiles</p> <p>2nd A1 for identifying the three outliers at 13, 87, 88 (dep on seeing both correct limits)</p>	
(d)	<p>M1 for drawing a box with only two whiskers one at each end</p> <p>1st A1ft for Q_1, Q_2 and Q_3 as a correctly drawn box (ft their values for Q_1, Q_2 and Q_3)</p> <p>2nd A1 for upper whisker ending at 76 (or 85) and lower whisker ending at 27 (or 21) NB</p> <p style="text-align: center;">Must be correctly paired.</p> <p>3rd A1 for the 3 outliers correctly shown (accuracy - half a small square throughout)</p> <p>SC fully correct but with both whiskers correct on each side. M0A1A0A1</p>	
(e)	<p>B1 for a correct comment, referring to ages, with reference to a correctly named statistic. Must include the figures compared.</p> <p>eg Females older than men and comparison of median, upper quartile or lower quartile, allow Q_1, Q_2 and Q_3 with their figures which must agree with the statement .</p> <p>eg Males ages more spread out than female and comparison of ranges with males = 75 and females = 73</p> <p>eg Females older than males since Males are symmetrical $[Q_3 - Q_2 : Q_2 - Q_1] 8 : 8$ Female are negative skew 5 : 11</p> <p>NB use of mean/ IQR/ minimum/ maximum is B0. Ignore incorrect comments.</p>	
(f)(i)	<p>B1 for deducing granddaughter is at or below lower quartile but not below 34</p> <p>Allow any reasonable adjustment for her mother's age, $\{34 \text{ to } x\}$ where $35 \leq x \leq 56$</p>	
(ii)	<p>M1 Suitable range for eldest daughter or Anja above upper quartile. Ignore any incorrect upper limit. May be implied by a correct range.</p> <p>A1 for a range of $[90\sim 93]$ for Anja's age</p>	

Question Number	Scheme	Marks										
<p>4. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p> $\text{Prob} = \frac{2}{3} \times \frac{1}{5} \times \frac{1}{8} + \frac{1}{3} \times \frac{4}{5} \times \frac{1}{8} + \frac{1}{3} \times \frac{1}{5} \times \frac{6}{8}$ $= \frac{1}{120} (2 + 4 + 6) \text{ or } (0.0166... + 0.033... + 0.05) = \frac{12}{120} \text{ or } \underline{\underline{\frac{1}{10}}}$ </p> <p> $P(RYY RYY \text{ or } YRY \text{ or } YYR) = \frac{\frac{2}{3} \times \frac{1}{5} \times \frac{1}{8}}{\frac{1}{10}} = \underline{\underline{\frac{1}{6}}}$ </p> <table border="1" data-bbox="277 1008 1292 1102"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>$P(X=x)$</td> <td>$\frac{64}{120}$ or $\frac{8}{15}$</td> <td>$\frac{42}{120}$ or $\frac{7}{20}$</td> <td>$\left[\frac{1}{10}\right]$</td> <td>$\frac{2}{120}$ or $\frac{1}{60}$</td> </tr> </table> <p> $E(X) = \frac{1}{120} (0 + "42" + "2 \times 12" + "3 \times 2") = \frac{72}{120} \text{ or } \underline{\underline{0.6}}$ </p>	x	0	1	2	3	$P(X=x)$	$\frac{64}{120}$ or $\frac{8}{15}$	$\frac{42}{120}$ or $\frac{7}{20}$	$\left[\frac{1}{10}\right]$	$\frac{2}{120}$ or $\frac{1}{60}$	<p>B1 B1 B1 (3)</p> <p>M1 A1ft A1*cso (3)</p> <p>M1 A1 (2)</p> <p>B1 M1 A1 (3)</p> <p>M1 A1 (2)</p> <p>[13]</p>
x	0	1	2	3								
$P(X=x)$	$\frac{64}{120}$ or $\frac{8}{15}$	$\frac{42}{120}$ or $\frac{7}{20}$	$\left[\frac{1}{10}\right]$	$\frac{2}{120}$ or $\frac{1}{60}$								
Notes												
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>1st B1 completing the structure of branches: 2, 4 then 7 or 8 and suitable labels e.g. Y or R'</p> <p>2nd B1 correct probabilities for at least bag A and bag B. Allow exact decimals</p> <p>3rd B1 for a fully correct tree diagram. Condone missing 0 as probability. Allow exact decimals</p> <p>1st M1 for at least one correct product of 3 probabilities (ft their tree diagram)</p> <p>1st A1ft for all 3 products of 3 probabilities added (no extras) (ft their tree diag.)</p> <p>2nd A1*cso for fully correct solution with no incorrect statements seen</p> <p>M1 for a ratio of probabilities with denominator of 0.1 and numerator $\frac{1}{60}$ oe or the product of 3 probabilities seen from their tree diagram representing $P(RYY)$ provided num < 0.1</p> <p>A1 for $\frac{1}{6}$ or exact equivalent</p> <p>B1 for a correct sample space i.e. $\{0, 1, 2, 3\}$ Allow extras if they have a probability of 0.</p> <p>M1 for at least 1 correct value of x and associated probability (excluding $x = 2$) [ft their tree]</p> <p>A1 for a fully correct probability distribution</p> <p>M1 for attempt at a correct expression (at least 2 correct ft part(d) non-zero products)</p> <p>A1 for 0.6 or any exact equivalent</p>											

Question Number	Scheme	Marks								
<p>5. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>[By symmetry $E(Y)] = \underline{0}$</p> <p>$q + r + u = \frac{19}{30}$</p> <p>$2(q + r) + u = 1$ [and attempt to solve e.g. $q + r = \dots$]</p> <p>$u = \frac{8}{30} = \frac{4}{15}$ (*)</p> <p>$E(Y^2) = (-9)^2 \times q + (-5)^2 \times r + 5^2 \times r + 9^2 q$ or $162q + 50r$</p> <p>$\text{Var}(Y) = 37 = E(Y^2) - "0"{}^2 \Rightarrow 37 = 162q + 50r$ oe</p> <p>Solving with $q + r = \frac{11}{30}$ oe e.g. $(162 - 50)q = 37 - \frac{55}{3}$ or</p> <p>$q = \frac{1}{6}$ and $r = \frac{1}{5}$</p> <p>$Y = 0 \Rightarrow D = 12, D = \sqrt{12^2 + Y^2}; Y = \pm 5 \Rightarrow D = 13$ or $Y = \pm 9 \Rightarrow D = 15$</p> <table border="1" data-bbox="277 965 1046 1104"> <tr> <td>d</td> <td>12</td> <td>13</td> <td>15</td> </tr> <tr> <td>$P(D = d)$</td> <td>$\frac{4}{15}$</td> <td>$\frac{6}{15}$ or $\frac{2}{5}$</td> <td>$\frac{5}{15}$ or $\frac{1}{3}$</td> </tr> </table>	d	12	13	15	$P(D = d)$	$\frac{4}{15}$	$\frac{6}{15}$ or $\frac{2}{5}$	$\frac{5}{15}$ or $\frac{1}{3}$	<p>B1</p> <p>(1)</p> <p>M1</p> <p>M1</p> <p>A1*cso</p> <p>(3)</p> <p>M1</p> <p>dM1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>B1, M1;A1</p> <p>M1A1ftA1ft</p> <p>(6)</p> <p>[14]</p>
d	12	13	15							
$P(D = d)$	$\frac{4}{15}$	$\frac{6}{15}$ or $\frac{2}{5}$	$\frac{5}{15}$ or $\frac{1}{3}$							
Notes										
	<p>(a) B1 for 0</p> <p>(b) 1st M1 for a correct equation in q, r and u using $F(0)$</p> <p>2nd M1 for a second equation clearly based on sum of probs = 1 and an attempt to solve these 2 equations</p> <p>A1* cso correct value for u found with no incorrect working</p> <p>(c) 1st M1 for an attempt at $E(Y^2)$ with at least 3 correct products seen. The negative numbers do not need to be in brackets</p> <p>2nd dM1 for attempt at correct equation in q and r using $\text{Var}(Y)$ [ft their $E(Y)$ and $E(Y^2)$] Condone missing subtraction of 0^2 if 0 in part(a)</p> <p>3rd M1 using $q + r = 11/30$ (awrt 0.37) to attempt to solve two linear equations in q and r leading to equation in one variable. May be implied by correct answers.</p> <p>A1 for $q = \frac{1}{6}$ and $r = \frac{1}{5}$ or exact equivalents</p> <p>(d) B1 for $D = 12$</p> <p>1st M1 for use of Pythagoras to work out $D = 13$ or 15</p> <p>1st A1 for $D = 13$ and 15</p> <p>2nd M1 for a correct value of D and an associated probability. Allow two occurrences (for 15 and 13) which add to the appropriate probability.</p> <p>2ndA1ft for two correct values of D and associated probs ft their +ve q and r if $q + r = \frac{11}{30}$ Allow two occurrences (for 15 and 13) which add to the appropriate probability.</p> <p>3rd A1ft for a fully correct probability distribution ft their +ve q and r if $q + r = \frac{11}{30}$</p>									

Question Number	Scheme	Marks
6. (a)	$H \sim N(25.1, 5.5^2)$ $P(H > 30.8) = P\left(Z > \frac{30.8 - 25.1}{5.5}\right)$ or $P(Z > 1.03636\dots)$ $= 1 - 0.8508$ $= 0.1492$ or better (calc: 0.1500...)	M1 M1 A1cso (3)
(b)	$[P(H < y) = 0.05 \text{ implies}] \quad \frac{y - 25.1}{5.5} = -1.6449$ $y = 16.053\dots$ so range is awrt 16.1 ~ 30.8	M1B1 A1 (3)
(c)(i)	$P(H < d) = 0.05 + 0.2 + 0.3 [= 0.55]$ $\frac{d - 25.1}{5.5} = 0.13$ (Calc 0.12566) $d = 0.13 \times 5.5 + 25.1 = 25.815$ (25.791... calc)	M1 M1 A1cso (3)
(ii)	$P(H < m) = 0.05 + 0.2 [= 0.25]$ $\frac{m - 25.1}{5.5} = -0.67$ (Calc 0.674489) $m = \text{awrt } \underline{21.4}$	M1 M1M1 A1 (4)
(d)	$\text{Height} = 2 \times "m" + 3 \times 25.8 + 3 \times 30.8 [+8]$ $= 220.6$ awrt 221 (cm)	M1 A1 (2)
[15]		
Notes		
(a)	1 st M1 for standardising 30.8 with 25.1 and 5.5 (allow \pm) Allow use of $z = 1.04$ 2 nd M1 for $1 - p$ (where $0.84 < p < 0.86$) A1cso for an answer of 0.1492... or better (calc: 0.1500...) with evidence of both M's seen	
(b)	M1 for standardising their letter y with 25.1 and 5.5 and setting equal to z value $1 < z < 2$ B1 for use of $z = \pm 1.6449$ or better (calc 1.6448536...) with the correct standardisation. A1 for awrt 16.1 (ISW)(calc 16.053305...) or range [16.1, 30.8](Allow $30.8 - 16.1 = 14.7[5]$) [awrt 16.05 scores 3/3 16.1 scores M1B0A1 unless 1.6449 or better is seen]	
Ans only		
(c)(i)	1 st M1 for a correct method to calculate $P(H < d)$ implied by $z = \text{awrt } 0.13$ Allow $0.05 + \text{awrt } 0.200 + \text{awrt } 0.300 [= 0.5505]$ 2 nd M1 for a correct standardisation $= z$ where $0.125 \text{ ,, } z \text{ ,, } 0.13$ A1cso both method marks awarded, no errors seen and awrt 25.82 or awrt 25.79 or $d = \text{awrt } 0.13 \times 5.5 + 25.1 = \text{awrt } 25.8$	
ALT	Verification 2 nd M1 allow $\frac{25.8 - 25.1}{5.5} = 0.127\dots$ or 0.13 A1 for 0.55 and 0.5517 (calc 0.5506 or better) seen	
(ii)	1 st M1 correct method for $P(H < m)$ Allow $0.05 + \text{awrt } 0.200$ implied by $ z = [0.67 - 0.68]$ 2 nd M1 for standardising m with 25.1 and 5.5 and setting equal to z value $(0.65 \square z \square 0.69)$ 3 rd M1 for standardising m with 25.1 and 5.5 and setting equal to awrt -0.67 oe A1 for $m = \text{awrt } 21.4$ (use of $z = 0.67$ gives 21.415... and $z = 0.68$ gives 21.36...) No need for 3 rd M1 to be awarded Answer only 21.4 gets M1M1M0A1. 21.39 gets 4/4	
(d)	M1 for $2 \times "m" + 3 \times 25.8 + 3 \times 30.8 [+n]$ where n is an integer $\square 0$ Allow m or ft their m A1 for awrt 221 (cm)	

Question Number	Scheme	Marks
ALT 1 (c)(i)	e.g. $P(H > 25.8 \mid "16.1" < H < 30.8)$ or $\frac{P(25.8 < H < 30.8)}{1 - (0.15 + 0.05)}$ $= \frac{0.8508 - 0.5517}{0.8}$ (tables) or $\frac{0.299345\dots}{0.8}$ (calc) $\approx \frac{3}{8}$	M1 M1 A1cso (3)
Notes		
(c)(i)	1 st M1 for a correct conditional probability statement ft their answer to (b) i.e. their y 2 nd M1 for a ratio of probs of the form $\frac{q}{0.8}$ where $q = 0.3$ to 1sf A1 for probability of approx $\frac{3}{8}$	