## Paper 1 (4CH1/1C and 4SD0/1C)

Question number	Answer	Mark
1(a)	Nucleus	1

Question number	Answer	Mark
1(b)	Proton	1

Question number	Answer	Additional guidance	Mark
1(c)	Equal numbers of protons and electrons	accept equal numbers of positive and negative particles/charges	1

Question number	Answer	Mark
1(d)	5	1

Question number	Answer	Mark
1(e)	Lithium	1

**Total for Question 1 = 5 marks** 

Question number	Answer	Additional guidance	Mark
2(a)(i)	An explanation that makes reference to the following two points:		
	<ul> <li>boxes 1 and 2 (1)</li> <li>because they both have only one type of atom/molecule (1)</li> </ul>	accept other indications, e.g. only He and only H-H accept species in place of atom/molecule	
		second mark can be awarded if only box 1 or box 2 identified	2

Question number	Answer	Additional guidance	Mark
2(a)(ii)	An explanation that makes reference to the following two points:  • boxes 3 and 5 (1)  • because they both have two different molecules (1)	second mark can be awarded if only box 3 or box 5 identified	2

Question number	Answer	Mark
2(b)(i)	Simple distillation	1

Question number	Answer	Mark
2(b)(ii)	Chromatography	1

Question number	Answer	Mark
2(b)(iii)	Crystallisation	1

**Total for Question 2 = 7 marks** 

Question number	Answer	Mark
3(a)	Reversible arrow	1

Question number	Answer	Additional guidance	Mark
3(b)(i)	<ul><li>(X) ammonium chloride (1)</li><li>(Y) ammonia and hydrogen chloride (1)</li></ul>	accept formulae	2

Question number	Answer	Mark
3(b)(ii)	D (subliming)	1

Question number	Answer	Additional guidance	Mark
3(c)	<ul> <li>An explanation that makes reference to the following three points:</li> <li>C (1)</li> <li>because ammonia molecules have lower mass or smaller M<sub>r</sub> (hence travel faster) (1)</li> <li>and so travel further in the same time (1)</li> </ul>	accept reverse arguments for hydrogen chloride	3

**Total for Question 3 = 7 marks** 

Question number	Answer	Additional guidance	Mark
4(a)	• 35 (1) • 41 (1)	final answer consequential on syringe readings	2

Question number	Answer	Additional guidance	Mark
4(b)	<ul> <li>Calculation of volume of oxygen used</li> <li>Calculation of original volume of air</li> <li>Calculation of percentage</li> <li>Example calculation: 80 - 43 = 37 (cm³) (1) 100 + 10 + 80 = 190 (cm³) (1) (37 × 100) ÷ 190 (= 19.47%) = 19% (1)</li> </ul>	accept 19.47% or 19.5%	2

question number	answer	mark
4(c)	<ul> <li>Decreased (1)</li> <li>Decreased (1)</li> <li>No effect (1)</li> </ul>	3

**Total for Question 4 = 8 marks** 

Question number	Answer	Mark
5(a)	<ul> <li>Any two of:</li> <li>concentration of copper(II) sulfate solution (1)</li> <li>volume of copper(II) sulfate solution (1)</li> <li>particle size of metal (1)</li> </ul>	2

Question number	Answer	Additional guidance	Mark
5(b)(i)	• (G) 5.5 (°C) (1) • (H) 11.5 (°C) (1)	accept 5.47	2

Question number	Answer	Mark
5(b)(ii)	<ul> <li>An explanation that makes reference to the following two points:</li> <li>H (1)</li> </ul>	
	because of the biggest temperature increase (1)	2

Question number	Answer	Additional guidance	Mark
5(b)(iii)	<ul> <li>An explanation that makes reference to the following two points:</li> <li>F (1)</li> <li>because there is no temperature increase (1)</li> </ul>	accept there is no reaction	
	, ,		2

**Total for Question 5 = 8 marks** 

Question number	Answer	Additional guidance	Mark
6(a)	The atoms of both elements have one electron in the outer shell	accept highest energy level in place of outer shell	1

Question number	Answer	Additional guidance	Mark
6(b)(i)	A description that makes reference to any two of the following points:	accept highest energy level in place of outer shell	
	<ul> <li>sodium floats/moves across the water (1)</li> <li>sodium melts (1)</li> <li>sodium disappears/gets smaller (1)</li> <li>effervescence/fizzing/bubbles/gas given off (1)</li> <li>white trail (1)</li> </ul>	accept forms a ball accept sodium dissolves ignore name of gas	2

Question number	Answer	Additional guidance	Mark
6(b)(ii)	An explanation that makes reference to the following points:		
	<ul><li> (final colour is) purple/blue (1)</li><li> because the solution is alkaline (1)</li></ul>	accept sodium hydroxide forms/ solution has high pH	
			2

Question number	Answer	Mark
6(b)(iii)	D (12)	1

Question number	Answer	Mark
6(c)	Lithium	1

Question number	Answer	Additional guidance	Mark
6(d)	Potassium catches fire	accept lilac/purple/violet flame	1

Question number	Answer	Additional guidance	Mark
6(e)	<b>2</b> Rb + <b>2</b> H <sub>2</sub> O → <b>2</b> RbOH + H <sub>2</sub> (1)	accept multiples and fractions	1

**Total for Question 6 = 9 marks** 

Question number	Answer	Mark
7(a)(i)	B and E	1

Question number	Answer	Mark
7(a)(ii)	(the only one that shows) All atoms and all bonds	1

Question number	Answer	Mark
7(a)(iii)	<ul> <li>D and F (1)</li> <li>they have the same molecular formula/the same number of each type of atom (1)</li> <li>but different structures/atoms joined together in different ways/different structural formulae (1)</li> </ul>	3

Question number	Answer	Mark
7(a)(iv)	4	1

Question number	Answer	Mark
7(b)(i)	<ul> <li>An explanation that makes reference to the following points:</li> <li>incomplete combustion/lack of oxygen (1)</li> <li>leads to the formation of carbon monoxide (1)</li> </ul>	2

Question number	Answer	Mark
7(b)(ii)	It reduces the capacity of blood to transport oxygen	1

Question number	Answer	Additional guidance	Mark
7(c)	<ul> <li>An explanation that makes reference to the following points:</li> <li>nitrogen in the air and oxygen react (1)</li> <li>at high temperatures (1)</li> <li>which causes the formation of nitrogen oxide(s) (1)</li> <li>oxides then react with water (vapour) in the atmosphere to form nitric acid (1)</li> </ul>	accept equation and formulae such as NO/NO <sub>2</sub> /NO <sub>x</sub> accept nitrous acid and formulae	4

**Total for Question 7 = 13 marks** 

Question number	Answer	Mark
8(a)	One reaction product is a gas and so escapes from the flask	1

Question number	Answer	Mark
8(b)(i)	<ul> <li>Any one of:</li> <li>balance reading recorded too late</li> <li>acid concentration greater than recorded</li> </ul>	
		1

Question number	Answer	Mark
8(b)(ii)	Loss in mass directly proportional to acid concentration	1

Question number	Answer	Additional guidance	Mark
8(c)	An explanation that makes reference to the following two points:		
	<ul> <li>more particles in the same volume <ul><li>(1)</li><li>so collide more frequently</li><li>(with malachite) (1)</li></ul> </li> </ul>	accept particles closer together	
			2

**Total for Question 8 = 5 marks** 

Question number	Answer	Mark
9(a)	<ul> <li>A description that makes reference to five of the following points:</li> <li>crude oil is heated/vaporised (1)</li> <li>the vapour enters the lower part of the column (1)</li> <li>there is a temperature gradient up the column (1)</li> <li>the vapour in the diesel fraction rises up the column until it condenses (1)</li> <li>at a height where its boiling point is lower than the temperature in the column (1)</li> <li>so the diesel fraction is removed (1)</li> </ul>	5

Question number	Answer	Additional guidance	Mark
9(b)	An explanation that makes reference to the following three points:		
	<ul> <li>dodecane contains hydrogen and carbon (1)</li> <li>only/and no other elements (1)</li> <li>and contains only single bonds (1)</li> </ul>	accept does not contain double bonds/multiple bonds	3

Question number	Answer	Mark
9(c)	С	1

Question number	Answer	Mark
9(d)	$C_8H_{18}$	1

Question number	Answer	Additional guidance	Mark
9(e)(i)	Ultraviolet radiation	accept ultraviolet light	1

Question number	Answer	Mark
9(e)(ii)	HCI	1

Question number	Answer	Additional guidance	Mark
9(e)(iii)	<ul> <li>All 6 atoms with a dot and cross representing each bonding pair of electrons (1)</li> <li>3 lone pairs of electrons on Cl and none on any of the H atoms (1)</li> </ul>	accept 2 dots or 2 crosses for each bond accept any combination of dots and crosses	
	atoms (1)		2

Question number	Answer	Mark
9(e)(iv)	Substitution	1

Question number	Answer	Mark
9(f)(i)	D	1

9(f)(ii) •	Dividing perce Dividing resul <b>OR</b>		omic masses (1	L)	
	obtaining rati Writing empir kample calculat	ical formula (			
	$C$ $25.9$ $12$ $2.16$ $3$ $gH_7BrO$ $ccept symbols in$	H 5.0 1 5.0 7 n any order	Br <u>57.6</u> 80 0.72 1	0 <u>11.5</u> 16 0.72 1	3

**Total for Question 9 = 19 marks** 

Question number	Answer	Mark
10(a)	<ul> <li>Increment in volume smaller/more precise (1)</li> <li>Avoids refilling the measuring cylinder (1)</li> </ul>	2

Question number	Answer		Additional guidance	Mark
10(b)				
	thermometer reading at end/°C	(26.8)	1 mark for temperature at start	
	thermometer reading at start/°C	18.7	1 mark for temperature rise consequential on readings	
	temperature rise/°C	8.1		2

Question number	Answer	Mark
<b>10</b> (c)(i)	29.5	1

Question number	Answer	Mark
10(c)(ii)	20.8	1

Question number	Answer	Mark
10(d)	Calculation of volume/mass of mixture	
	Calculation of temperature increase	
	• Substitution of values into $q=mc\Delta T$	
	Calculation of heat energy released with unit	
	Example calculation:	
	$20.0 + 20.0 = 40.0 \text{ (cm}^3\text{) (1)}$	
	$30.0-18.5 = 11.5 (^{\circ}C) (1)$	
	$q = 40.0 \times 4.2 \times 11.5$ (1)	
	$q = 1900 \mathrm{J} (1) (\mathrm{accept}  1932 \mathrm{J})$	
		4

Question number	Answer	Mark
10(e)	<ul> <li>Setting out of ΔH calculation</li> <li>Division by 1000 to obtain answer in kJ/mol</li> <li>Example calculation:</li> <li>1600 ÷ 0.040 (1)</li> <li>= -40 (kJ/mol) (1)</li> </ul>	2

Total for Question 10 = 12 marks

Question number	Answer		Mark
11(a)	1 mark for each box completed correct	У	
	Reactants Name of salt formed	Other product(s)	
	(zinc + hydrochloric zinc chloride acid)	hydrogen	
	(calcium carbonate + nitric acid)	water + carbon dioxide	4

Question number	Answer	Mark
11(b)(i)	<ul><li>Use excess aluminium hydroxide (1)</li><li>Stir (thoroughly) (1)</li></ul>	2

Question number	Answer	Mark
11(b)(ii)	To remove unreacted aluminium hydroxide/solid	1

Question number	Answer	Mark
11(b)(iii)	<ul> <li>Any one of:</li> <li>leave in a warm place (1)</li> <li>use filter paper or paper towel (1)</li> </ul>	1

Question number	Answer	Mark
11(c)	<ul> <li>Calculation of M<sub>r</sub> of aluminium hydroxide</li> <li>Calculation of amount of aluminium hydroxide</li> <li>Reference to 2: 3 ratio in equation AND statement that sulfuric acid is in excess</li> </ul>	
	Example calculation: $27 + (3 \times 17) = 78 (1)$ $3.9 \div 78 = 0.05 \text{ mol } (1)$	
	This is more than 3/2 times amount of aluminium hydroxide, so sulfuric acid is in excess (1) (accept other valid methods of calculation)	3

Question number	Answer	Mark
11(d)	Calculation of $M_r$ of aluminium sulfate setting out calculation of mass final answer  Example calculation: $(27 \times 2) + (32 \times 3) + (16 \times 12) = 342(1)$	
	mass = 342 x 0.25 (1) 85.5 g (1)	3

Question number	Answer	Additional guidance	Mark
11(e)	<ul> <li>Calculation of amount of lead(II) nitrate</li> <li>Percentage method</li> <li>Percentage answer</li> <li>Example calculation:</li> <li>209 ÷ 331 = 0.631 mol (1)</li> <li>0.631 x 100 (1) = 84% (1)</li> <li>0.75</li> </ul>	allow full credit for calculations using masses	
	3.73		3

Total for Question 11 = 17 marks

**TOTAL FOR PAPER = 110 MARKS**